



GOLDMAN ALGEBRA, OPERS AND THE SWAPPING ALGEBRA

François Labourie

► To cite this version:

François Labourie. GOLDMAN ALGEBRA, OPERS AND THE SWAPPING ALGEBRA. 2016.
hal-01329439

HAL Id: hal-01329439

<https://hal.univ-cotedazur.fr/hal-01329439>

Preprint submitted on 9 Jun 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

GOLDMAN ALGEBRA, OPERS AND THE SWAPPING ALGEBRA

FRANÇOIS LABOURIE

ABSTRACT. We define a Poisson Algebra called the *swapping algebra* using the intersection of curves in the disk. We interpret a subalgebra of the fraction algebra of the swapping algebra – called the *algebra of multifractions* – as an algebra of functions on the space of cross ratios and thus as an algebra of functions on the Hitchin component as well as on the space of $\mathrm{SL}_n(\mathbb{R})$ -opers with trivial holonomy. We relate this Poisson algebra to the Atiyah–Bott–Goldman symplectic structure and to the Drinfel’d–Sokolov reduction. We also prove an extension of Wolpert formula.

1. INTRODUCTION

The purpose of this article is threefold. We first introduce the *swapping algebra* which is a Poisson algebra generated – as a commutative algebra – by pairs of points on the circle. Then we relate this construction to two well known Poisson structures:

- the Poisson structure of the character variety of representations of a surface group in $\mathrm{PSL}_n(\mathbb{R})$ discovered by Atiyah, Bott and Goldman [1, 6, 7]
- the Poisson structure of the space of $\mathrm{PSL}_n(\mathbb{R})$ -opers introduced by Dickey, Gel’fand and Magri and described in a geometrical way by Drinfel’d and Sokolov [18, 4, 3].

One way to interpret heuristically these relations is to say that the swapping algebra embodies the notion of a “Poisson structure” for the space of all cross ratios, space that contains both the space of opers and the “universal (in genus) Hitchin component”. As a byproduct of the methods of this paper, we also produce a generalisation of the Wolpert formula which computes the brackets of length functions for the Hitchin component.

Univ. Paris-Sud, Laboratoire de Mathématiques, Orsay F-91405 Cedex; CNRS, Orsay cedex, F-91405. The research leading to these results has received funding from the European Research Council under the *European Community’s* seventh Framework Programme (FP7/2007-2013)/ERC *grant agreement* n° FP7-246918, as well as well as from the ANR program ETTT (ANR-09-BLAN-0116-01) .

The results of this article were announced in [16]. The relation – at a topological level – between the character variety and opers was already noted by the author in [13], by Fock and Goncharov in [5] and foreseen by Witten in [24] (see also [9],[8]). I thank Martin Bridson, Louis Funar and Bill Goldman and for their interest and help.

We now explain more precisely the content of this article.

1.1. The swapping algebra. Our first result is the construction of the *swapping algebra*. To avoid cumbersome expressions, we shall denote most of the time the ordered pair (X, x) of points of the circle by the concatenated symbol Xx . We recall in Paragraph 2.1 the definition and properties of the linking number $[Xx, Yy]$ of the two pairs (X, x) and (Y, y) . If P is a subset of the circle, we denote by $\mathcal{L}(P)$ the commutative associative algebra generated by pairs of points of P with the relations $XX = 0$, for all X in P . Our starting result is the following

Theorem 1. [SWAPPING BRACKET] *For every number α , there exists a unique Poisson bracket on $\mathcal{L}(P)$ such that the bracket of two generators is*

$$\{Xx, Yy\}_\alpha = [Xx, Yy](Xy.Yx + \alpha.Xx.Yy).$$

The *swapping algebra* is the algebra $\mathcal{L}_\alpha(P)$ endowed with the Poisson bracket $\{, \}_\alpha$. This theorem is proved in Section 2. The goal of this paper is to relate this swapping algebra to other Poisson algebras.

One should note that this bracket can be used to express very simply some results of Wolpert and in particular the variation of the length of curve transverse to a shear [26, 25].

1.2. Cross ratios and the multi fraction algebra. We shall concentrate on the interpretation of an offshoot of the swapping algebra. We denote by $\mathcal{Q}_\alpha(P)$ the algebra of fractions of $\mathcal{L}_\alpha(P)$ equipped with the induced Poisson structure. The *multi fraction algebra* $\mathcal{B}(P)$ is the vector subspace of $\mathcal{Q}_\alpha(P)$ generated by the *elementary multi fractions*:

$$[X, x; \sigma] := \frac{\prod_{i=1}^{i=n} X_i x_{\sigma(i)}}{\prod_{i=1}^{i=n} X_i x_i},$$

where $X = (X_1, \dots, X_n)$ and $x = (x_1, \dots, x_n)$ are n t -uples of points of P and σ is a permutation of $\{1, \dots, n\}$. Then we have the easy proposition

Proposition 2. *The multi fraction algebra is a Poisson subalgebra of $\mathcal{Q}_\alpha(P)$. The induced Poisson structure does not depend on α . Finally*

$\mathcal{B}(\mathbf{P})$ is generated as a commutative algebra by the cross fractions:

$$[X, Y, x, y] := \frac{Xx.Yy}{Yx.Xy}.$$

In particular, it follows that the multi fraction algebra is naturally mapped to the commutative algebra of functions on cross ratios (See Section 3). Thus the existence of a Poisson structure on the multi fraction algebra can be interpreted as that of a Poisson structure on the space of cross ratios.

1.3. The multi fraction algebra as a “universal” Goldman algebra. We then relate the multi fraction algebra to the Goldman algebra. Let Γ be the fundamental group of a surface S and \mathbf{P} be the subset of $\partial_\infty \Gamma$ consisting of fixed points of elements of Γ . The Hitchin component $\mathbf{H}(n, S)$ of the character variety of representations of Γ in $\mathrm{PSL}_n(\mathbb{R})$ was interpreted in [14] as a space of cross ratios. Thus every multi fraction in $\mathcal{B}(\mathbf{P})$ gives a smooth function on the Hitchin component (see Proposition 4.2.4 for details). Thus we have a restriction

$$l_S : \mathcal{B}(\mathbf{P}) \rightarrow C^\infty(\mathbf{H}(n, S)).$$

This mapping is not a Poisson morphism, nevertheless it becomes so when we take sequences of well chosen finite index subgroups. More precisely, we define and prove, as an immediate consequence of one of the main result of Niblo in [19], the existence of *vanishing sequences* of finite index subgroups $\{\Gamma_n\}_{n \in \mathbb{N}}$ of Γ ; these sequences are essentially such that every geodesic becomes eventually simple and for which the intersection of two geodesics becomes eventually minimal (See Paragraph 6.2.1 and Appendix 11 for precisions).

Then denoting by $\{, \}_W$ the swapping bracket and $\{, \}_{S_n}$ the Goldman bracket for $S_n := \tilde{S}/\Gamma_n$ coming from the Atiyah–Bott–Goldman symplectic form on the character variety, we prove in Section 9

Theorem 3. [GOLDMAN BRACKET FOR VANISHING SEQUENCES]

Let $\{\Gamma_m\}_{m \in \mathbb{N}}$ be a vanishing sequence of subgroups of $\pi_1(S)$. Let $\mathbf{P} \subset \partial_\infty \pi_1(S)$ be the set of end points of geodesics. Let b_0 and b_1 be two multi fractions in $\mathcal{B}(\mathbf{P})$. Then we have,

$$\lim_{n \rightarrow \infty} \{b_0, b_1\}_{S_n} = \{b_0, b_1\}_W. \quad (1)$$

The statement of this theorem actually requires some preliminaries in defining properly the meaning of Assertion (1). In a way, this result tells us that the swapping bracket is the Goldman bracket on the universal solenoid.

The proof relies on the description of special multi fractions called *elementary functions* (see Paragraph 4.2) as limits of the well studied functions on the character variety known as *Wilson loops*.

Another result is a precise asymptotic formula, on a fixed surface this time, relating the Goldman and the swapping brackets. Let Γ be as above. Let $\gamma \in \Gamma$. Let finally $y \in \mathbb{P}$, and γ^+ , γ^- be respectively the attractive and repulsive fixed points of γ in $\partial_\infty(\Gamma)$. Let us define the following formal series of cross fractions

$$\hat{\ell}_\gamma(y) = \log \left(\frac{\gamma^+ \gamma^{-1}(y) \cdot \gamma^- \gamma(y)}{\gamma^+ \gamma(y) \cdot \gamma^- \gamma^{-1}(y)} \right)$$

In [14], we show that the *period function* $\ell_\gamma := I_S(\hat{\ell}_\gamma(y))$ – seen as a function on the character variety – is independent on y and is a function of the eigenvalues of the monodromy of γ . These period functions coincide with the length functions for the classical Teichmüller Theory – that is $n = 2$.

We now have

Theorem 4. [BRACKET OF LENGTH FUNCTIONS] *Let γ and η be homotopy classes of curves which as simple curves have at most one intersection point, then we have*

$$\lim_{n \rightarrow \infty} I_S(\{\hat{\ell}_{\gamma^n}(y), \hat{\ell}_{\eta^n}(y)\}_W) = \{\ell_\gamma, \ell_\eta\}_S$$

As a tool of the proof of this result we prove the following extension of the Wolpert formula [26, 25].

Theorem 5. [GENERALISED WOLPERT FORMULA] *Let γ and η be two homotopy classes of curves which as simple curves have exactly one intersection point. Then the Goldman bracket of the two length functions ℓ_γ and ℓ_η is*

$$\{\ell_\gamma, \ell_\eta\}_S(\mathbf{b}) = \iota(\gamma, \eta) \sum_{\varepsilon, \varepsilon' \in \{-1, 1\}} \varepsilon \cdot \varepsilon' \cdot \mathbf{b} \left(\gamma^\varepsilon, \eta^{\varepsilon'}, \gamma^{-\varepsilon}, \eta^{-\varepsilon'} \right). \quad (2)$$

1.4. The multi fraction algebra and $\mathrm{PSL}_n(\mathbb{R})$ -opers. We finally relate the multi fraction algebra to opers. We recall in Section 10 the definition of real opers and their interpretation as maps to the projective space $\mathbb{P}(\mathbb{R}^n)$ and its dual. In particular, opers with trivial holonomy can be embedded in the space of smooth cross ratios. The Drinfel'd-Sokolov reduction allows us to define the Poisson bracket of pairs of *acceptable observables*, a subclass of functions on the spaces of opers. We then show that this Poisson bracket coincide with the swapping bracket

Theorem 6. [SWAPPING BRACKET AND OPERS]

Let $(X_0, x_0, Y_0, y_0, X_1, x_1, Y_1, y_1)$ be pairwise distinct points on the circle \mathbb{T} . Then the cross fractions $[X_0, x_0, Y_0, y_0]$ and $[X_1, x_1, Y_1, y_1]$ defines a pair of acceptable observables whose Poisson bracket with respect to the Drinfel'd-Sokolov reduction coincide with their Poisson bracket in the multi fraction algebra.

CONTENTS

1. Introduction	1
1.1. The swapping algebra	2
1.2. Cross ratios and the multi fraction algebra	2
1.3. The multi fraction algebra as a “universal” Goldman algebra	3
1.4. The multi fraction algebra and $\mathrm{PSL}_n(\mathbb{R})$ -opers	4
2. The swapping bracket	6
2.1. Linking number for pairs of points	6
2.2. The swapping algebra	12
2.3. The multi fraction algebra	15
3. Cross ratios and cross fractions	16
3.1. Cross ratios	16
3.2. Multi fractions as functions	17
3.3. Multi fractions and Hitchin components	17
4. Wilson loops, multi fractions and length functions	19
4.1. Wilson loops	19
4.2. Elementary functions	21
4.3. The swapping bracket of elementary functions	23
4.4. Length functions	24
5. The Goldman algebra	25
5.1. The Atiyah-Bott-Goldman symplectic form	25
5.2. Wilson loops and the Goldman algebra	26
6. Vanishing sequences and the main results	27
6.1. Poisson brackets of length functions	27
6.2. Poisson brackets of multi fractions	28
7. Product formulas and bouquet in good position	29
7.1. An alternative formulation of the Goldman bracket	30
7.2. The product formula	31
7.3. Bouquets in good position and the product formula	33
7.4. Bouquets and covering	37
7.5. Finding bouquets in good position	37
8. Asymptotics	42
8.1. Properties of vanishing sequences	43

8.2.	Asymptotic product formula for Wilson loops	47
8.3.	Asymptotics of brackets of multifractions	55
9.	Goldman and swapping algebras: proofs of the main results	57
9.1.	Poisson brackets of elementary functions and the proof of Theorem 6.2.4	58
9.2.	Poisson brackets of length functions	59
10.	Drinfel'd-Sokolov reduction	61
10.1.	Opers and non slipping connections	61
10.2.	The Poisson structure on the space of connections	64
10.3.	Drinfel'd-Sokolov reduction	66
10.4.	Opers and Frenet curves	68
10.5.	Cross ratios and opers	72
10.6.	Poisson brackets on the space of connections	74
10.7.	Drinfel'd-Sokolov reduction and the multi fraction algebra	77
11.	Appendix: existence of vanishing sequences	78
	References	79

2. THE SWAPPING BRACKET

In this section, we first recall the properties and definition of the linking number of two ordered pairs of points. We then construct the swapping algebra and prove Theorem 1 which relies on an identity involving the linking numbers of six points.

2.1. Linking number for pairs of points. We recall that if (X, x, Y, y) is a quadruple of points on the real line the *linking number* of (X, x) and (Y, y) is

$$[Xx, Yy] := \frac{1}{2} \left(\text{Sign}(X - x) \text{Sign}(X - y) \text{Sign}(y - x) - \text{Sign}(X - x) \text{Sign}(X - Y) \text{Sign}(Y - x) \right), \quad (3)$$

where $\text{Sign}(x) = -1, 0, 1$ whenever $x < 0, x = 0$ and $x > 0$ respectively. By definition, the linking number is invariant by orientation preserving homeomorphisms of the real line.

- (1) When the four points are pairwise distinct, this linking number is also the total linking number of the curve joining X to x with the curve joining Y to y in the upper half plane.
- (2) The equality cases are as follows:

(a) For all points (X, Y, y) on the circle

$$[XX, Yy] = 0 = [Xy, Xy]. \quad (4)$$

(b) If, up to cyclic permutation, (X, Y, x) are pairwise points and oriented, then

$$[Xx, Yx] = 1/2 \quad (5)$$

The first observation shows that we can define the linking number of a quadruple of points on the oriented circle S^1 by choosing a point x_0 disjoint from the quadruple and defining the linking number as the linking number of the quadruple in $S^1 \setminus \{x_0\} \sim \mathbb{R}$. The linking number so defined does not depend on the choice of x_0 and is invariant under orientation preserving homeomorphisms.

2.1.1. *Properties of the linking number.* We abstract the useful property (for us) of the linking number of pairs of points in the following definition. Let P be any set.

Definition 2.1.1. *A linking number on pair of points of P is a map from P^4 to a commutative ring*

$$(X, x, Y, y) \rightarrow [Xx, Yy],$$

so that for all points X, x, Y, y, Z, z

$$[Xx, Yy] + [Yy, Xx] = 0, \quad \text{FIRST ANTISYMMETRY,} \quad (6)$$

$$[Xx, Yy] + [Xx, yY] = 0, \quad \text{SECOND ANTISYMMETRY,} \quad (7)$$

$$[zy, XY] + [zy, YZ] + [zy, ZX] = 0, \quad \text{COCYCLE IDENTITY,} \quad (8)$$

and moreover if (X, x, Y, y) are all pairwise distinct then

$$[Xx, Yy] \cdot [Xy, Yx] = 0, \quad \text{LINKING NUMBER ALTERNATIVE.} \quad (9)$$

We illustrate the cocycle identity and the alternative for the standard linking number in Figure (1)

Then we prove

Proposition 2.1.2. *The canonical linking number on pair of points of the circle is a linking number in the sense of the previous definition.*




FIGURE 1. linking number for pairs of points on the circle

Proof. The first two symmetries are checked from the definition. When $\{x, y\} \cap \{X, Y, Z\} = \emptyset$, then Equation (8) follows from the geometric definition of the linking number. It remains to check different cases of equality. We can assume that (X, Y, Z) are pairwise distinct: otherwise the equality follows from the two previous ones and (4).

- if $x = y$, the equation is true by (4).
- Assume that up to cyclic permutations of (X, Y, Z) we have $x = X$ and $y \notin \{X, Y, Z\}$, then the equality follows from the following remark. Let z, t be points close enough to x so that (z, x, t) is oriented then when $A = Y$ or $A = Z$, we have

$$[xy, xA] = \frac{1}{2} ([zy, xA] + [ty, xA]).$$

- Assume finally that $(x, y) = (X, Y)$, the the equality reduces to

$$[XY, ZX] + [XY, YZ] = 0,$$

which is true, by Equation (5) and the fact that (X, Y, Z) has the opposite orientation of (Y, X, Z) .

Equation (9) follows from the geometric definition of linking number. \square

A linking number satisfies more complicated relations. Namely

Proposition 2.1.3. *Let (X, x, Z, z, Y, y) be 6 points on the set \mathbf{P} equipped with an intersection $[\cdot, \cdot]$, then*

$$[Xy, Zz] + [Yx, Zz] = [Xx, Zz] + [Yy, Zz]. \quad (10)$$

Moreover, if

$$\{X, x\} \cap \{Y, y\} \cap \{Z, z\} = \emptyset,$$

then

$$[Xx, Yy][Xy, Zz] + [Zz, Xx][Zx, Yy] + [Yy, Zz][Yz, Xx] = 0 , \quad (11)$$

$$[Xx, Yy][Yx, Zz] + [Zz, Xx][Xz, Yy] + [Yy, Zz][Zy, Xx] = 0 . \quad (12)$$

REMARKS:

- (1) We remark that the hypothesis on the configuration of points is necessary: if X, x, Y, Z are pairwise distinct, then $F(X, x, Y, x, Z, x) \neq 0$ in the case of the standard linking number on pair of points on the circle.
- (2) A simple way to prove this proposition is to use a mathematical computing software, we give below a mathematical proof.

Proof. Formula (10) follows at once from the cocycle identity (8). We now prove Formulas (11) and (12). Let us define

$$F(X, x, Y, y, Z, z) := [Xx, Yy][Xy, Zz] + [Zz, Xx][Zx, Yy] + [Yy, Zz][Yz, Xx] ,$$

$$G(X, x, Y, y, Z, z) := [Xx, Yy][Yx, Zz] + [Zz, Xx][Xz, Yy] + [Yy, Zz][Zy, Xx] .$$

We first prove some symmetries of F and G .

Our first observation is that, using (6), we get that

$$F(X, x, Y, y, Z, z) = -G(Y, y, X, x, Z, z). \quad (13)$$

Thus we only need to prove that $F = 0$.

STEP 1: *The expression F is invariant under all permutations of the pairs $(X, x), (Y, y)$ and (Z, z)*

Using Equations (10) and (6), we obtain that

$$\begin{aligned} F(X, x, Y, y, Z, z) + G(X, x, Y, y, Z, z) &= [Xx, Yy][Yy, Zz] + [Xx, Yy][Xx, Zz] \\ &\quad + [Zz, Xx][Zz, Yy] + [Zz, Xx][Xx, Yy] \\ &\quad + [Yy, Zz][Yy, Xx] + [Yy, Zz][Zz, Xx] \\ &= 0 . \end{aligned}$$

Hence, by Equation (13).

$$F(X, x, Y, y, Z, z) = F(Y, y, X, x, Z, z) \quad (14)$$

By construction F is invariant by cyclic permutations and thus from the previous equation F is invariant by all permutations of the pairs $(X, x), (Y, y)$ and (Z, z) .

STEP 2: *The expression F satisfies a cocycle equation*

$$F(X, x, Y, y, Z, z) + F(x, t, Y, y, Z, z) = F(X, t, Y, y, Z, z). \quad (15)$$

We also have the symmetries

$$\begin{aligned}
F(X, x, Y, y, Z, z) &= -F(x, X, Y, y, Z, z) \\
&= -F(X, x, y, Y, Z, z) \\
&= -F(X, x, Y, y, z, Z).
\end{aligned} \tag{16}$$

The Symmetries (16) follow at once from the Cocycle (15) and the fact that $F(X, X, Y, y, Z, z) = 0$.

Let us prove a cocycle equation for F . We shall only use the cocycle Equation (8) and the previous symmetries for the linking number. By definition,

$$\begin{aligned}
F(X, x, Y, y, Z, z) + F(x, t, Y, y, Z, z) \\
&= [Xx, Yy][Xy, Zz] + [xt, Yy][xy, Zz] \\
&\quad + [Zz, Xx][Zx, Yy] + [Zz, xt][Zt, Yy] \\
&\quad + [Yy, Zz][Yz, Xx] + [Yy, Zz][Yz, xt].
\end{aligned}$$

Using Equation (8) to expand the first and regrouping the fifth and sixth terms of the righthand side, we get

$$\begin{aligned}
F(X, x, Y, y, Z, z) + F(x, t, Y, y, Z, z) \\
&= [Xt, Yy][Xy, Zz] + [tx, Yy][Xy, Zz] + [xt, Yy][xy, Zz] \\
&\quad + [Zz, Xx][Zx, Yy] + [Zz, xt][Zt, Yy] \\
&\quad + [Yy, Zz][Yz, Xt].
\end{aligned}$$

Using Equation (8) for regrouping the second and third term of the righthand side and rearranging, we get

$$\begin{aligned}
F(X, x, Y, y, Z, z) + F(x, t, Y, y, Z, z) \\
&= [Xt, Yy][Xy, Zz] \\
&\quad + [Zz, Xx][Zx, Yy] + [Zz, Xx][xt, Yy] + [Zz, xt][Zt, Yy] \\
&\quad + [Yy, Zz][Yz, Xt] \\
&= F(X, t, Y, y, Z, z).
\end{aligned}$$

Using Equation (8) to regroup the second and third term, then the fourth, of the right hand side, we finally get

$$\begin{aligned}
F(X, x, Y, y, Z, z) + F(x, t, Y, y, Z, z) \\
&= [Xt, Yy][Xy, Zz] \\
&\quad + [Zz, Xt][Zt, Yy] \\
&\quad + [Yy, Zz][Yz, Xt] \\
&= F(X, t, Y, y, Z, z).
\end{aligned}$$

STEP 4: *If (X, x, Y, y) are pairwise distinct, then*

$$F(X, x, Y, y, Y, x) = 0 \quad (17)$$

$$F(X, x, X, x, Y, y) = 0. \quad (18)$$

For Equation (17), it follows from the Alternative (9) and the cocycle Equation (8) that

$$\begin{aligned}
F(X, x, Y, y, Y, x) \\
&= [Xx, Yy][Xy, Yx] + ([Yx, Xx][Yx, Yy] + [Yy, Yx][Yx, Xx]) \\
&= 0.
\end{aligned}$$

This proves Formula (17). Similarly, using the cocycle Formula for F (15) for the first equality, symmetries for the second and our previous Formula (17) (for (X, x, y, Y)) for the last, we get

$$\begin{aligned}
F(X, x, X, x, Y, y) \\
&= F(X, x, X, y, Y, y) + F(X, x, y, x, Y, y) \\
&= F(X, x, y, x, y, Y) = F(X, x, y, Y, y, x) \\
&= 0.
\end{aligned}$$

STEP 4: *If (X, x, Y, y, Z) are pairwise distinct, then*

$$F(X, x, Y, y, Z, x) = 0 \quad . \quad (19)$$

Using the cocycle equation (15) for F and the previous step, we get

$$\begin{aligned}
F(X, x, Y, y, Z, x) \\
&= F(X, x, Y, Z, Z, x) + F(X, x, Z, y, Z, x) \\
&= -F(X, x, Z, Y, Z, x) = 0.
\end{aligned}$$

FINAL STEP: *If (X, x, Y, y, Z, z) are pairwise distinct, then*

$$F(X, x, Y, y, Z, z) = 0 \quad . \quad (20)$$

Indeed, we have using the cocycle Equation for F (15) for the first equality, symmetries for the second, on the previous step for the last, we get

$$\begin{aligned} F(X, x, Y, y, Z, z) &= F(X, x, Y, y, Z, Y) + F(X, x, Y, y, Y, z) \\ &= F(y, Y, x, X, Z, Y) - F(y, Y, x, X, y, z, Y) \\ &= 0 \quad . \end{aligned}$$

This concludes the proof \square

2.2. The swapping algebra. Let P be a set and $[\cdot, \cdot]$ be a linking number with values in an integral domain A . We represent a pair (X, x) of points of P by the expression Xx . We consider the free associative commutative algebra $\mathcal{L}(P)$ generated over A by pair of points on P , together with the relation $Xx = 0$.

Let α be any element in A . We define the *swapping bracket* of two pairs of points as the following element of $\mathcal{L}(P)$

$$\{Xx, Yy\}_\alpha := [Xx, Yy](\alpha.Xx.Yy + Xy.Yx). \quad (21)$$

We extend the swapping bracket to the whole algebra $\mathcal{L}(P)$ using the Leibniz Rule and call the resulting algebra $\mathcal{L}_\alpha(P)$ the *swapping algebra*.

Theorem 2.2.1. *The swapping bracket satisfies the Jacobi identity. Hence, the swapping algebra $\mathcal{L}_\alpha(P)$ is a Poisson algebra.*

Proof. All we need to check is the Jacobi identity

$$\{\{Xx, Yy\}_\alpha, Xz\}_\alpha + \{\{Xy, Xz\}_\alpha, Xx\}_\alpha + \{\{Xz, Xx\}_\alpha, Yy\}_\alpha = 0,$$

for the generators of the algebra.

We make preliminary computations, omitting the subscript α in the bracket. The triple bracket $\{\{A, B\}, C\}$ is a polynomial of degree 2 in α and we wish to compute its coefficients. By definition, using the Leibniz rule for the second equality, we have

$$\begin{aligned} \{\{Xx, Yy\}, Zz\} &= [Xx, Yy](\alpha\{Xx.Yy, Zz\} + \{Xy.Yx, Zz\}) \\ &= \alpha[Xx, Yy](\{Xx, Zz\}.Yy + \{Yy, Zz\}.Xx) \\ &\quad + [Xx, Yy](\{Xy, Zz\}.Yx + \{Yx, Zz\}.Xy). \quad (22) \end{aligned}$$

Now we compute two expressions appearing in the right hand side of the previous equation. We have

$$\begin{aligned} & \{Xx, Zz\}.Yy + \{Yy, Zz\}.Xx \\ &= \alpha ([Xx, Zz] + [Yy, Zz]) Xx.Yy.Zz \\ & \quad + ([Xx, Zz]Xz.Yy.Zx + [Yy, Zz]Xx.Yz.Zy) . \end{aligned} \quad (23)$$

Similarly

$$\begin{aligned} & \{Xy, Zz\}.Yx + \{Yx, Zz\}.Xy \\ &= \alpha ([Xy, Zz] + [Yx, Zz]) Xy.Yx.Zz \\ & \quad + [Xy, Zz]Xz.Yx.Zy + [Yx, Zz]Xy.Yz.Zx. \end{aligned} \quad (24)$$

It follows from Equations (23) and (24) that the coefficient of α^2 in the triple bracket (22) is

$$P_2 := ([Xx, Yy][Xx, Zz] + [Xx, Yy][Yy, Zz]) Xx.Yy.Zz. \quad (25)$$

The coefficient of α in the triple bracket (22) is

$$\begin{aligned} P_1 := & [Xx, Yy][Xx, Zz]Xz.Yy.Zx + [Xx, Yy][Yy, Zz]Xx.Yz.Zy \\ & + ([Xx, Yy][Xy, Zz] + [Xx, Yy][Yx, Zz])Xy.Yx.Zz. \end{aligned} \quad (26)$$

Finally the constant coefficient is

$$P_0 := [Xx, Yy][Xy, Zz]Xz.Yx.Zy + [Xx, Yy][Yx, Zz]Xy.Yz.Zx, \quad (27)$$

so that

$$\{\{Xx, Yy\}, Zz\} = \alpha^2 P_2 + \alpha P_1 + P_0. \quad (28)$$

In order to check the Jacobi identity, we have to consider the sum S_2 , S_1 and S_0 over cyclic permutations of (Xx, Yy, Zz) of the three terms P_2 , P_1 and P_0 . We consider successivley these three coefficients.

TERM OF DEGREE 0: We first have

$$S_0 = F(X, x, Y, y, Z, z)(Xz.Yx.Zy - Xy.Yz.Zx), \quad (29)$$

Indeed, we have

$$S_0 = A.Xz.Yx.Zy + B.Xy.Yz.Zx,$$

where

$$\begin{aligned} A &= [Xx, Yy][Xy, Zz] + [Zz, Xx][Zx, Yy] + [Yy, Zz][Yz, Xx] = F(X, x, Y, y, Z, z), \\ B &= [Xx, Yy][Yx, Zz] + [Zz, Xx][Xz, Yy] + [Yy, Zz][Zy, Xx] = G(X, x, Y, y, Z, z). \end{aligned}$$

Now Equation (29) follows from Equation (13).

It follows from Proposition ?? that if

$$\{X, x\} \cap \{Y, y\} \cap \{Z, z\} = \emptyset,$$

then $F = 0$, hence $S_0 = 0$.

Up to cyclic permutations, we just have to consider two cases

- (1) If $x = y = z$ or $X = Y = Z$ then

$$Xz.Yx.Zy - Xy.Yz.Zx = 0,$$

hence $S_0 = 0$,

- (2) If $x = y = Z$ or $X = Y = z$ or the other cases obtained by cyclic permutations, since $aa = 0$, we have

$$Xz.Yx.Zy = Xy.Yz.Zx = 0$$

Thus $S_0 = 0$.

We have completed the proof that $S_0 = 0$.

TERM OF DEGREE 1. Next, we write

$$\begin{aligned} P_1 &= A_1(X, x, Y, y, Z, z)Xx.Yz.Zy \\ &\quad + A_2(X, x, Y, y, Z, z)Xz.Yy.Zx \\ &\quad + A_3(X, x, Y, y, Z, z)Xy.Yx.Zz . \end{aligned}$$

Thus

$$S_1 = A_x.Xx.Yz.Zx + A_y.Xz.Yy.Zx + A_z.Xy.Yx.Zz ,$$

where

$$\begin{aligned} A_z &= A_3(X, x, Y, y, Z, z) + A_2(Y, y, Z, z, X, x) + A_1(Z, z, X, x, Y, y) \\ &= [Xx, Yy][Xy, Zz] + [Xx, Yy][Yx, Zz] \\ &\quad + [Yy, Zz][Yy, Xx] + [Zz, Xx][Xx, Yy] \\ &= [Xx, Yy] ([Xy, Zz] + [Yx, Zz] - [Yy, Zz] - [Xx, Zz]) . \end{aligned}$$

By Equation (10), $A_z = 0$. Therefore, $A_y = A_z = A_x = 0$ by cyclic permutations. We have completed the proof that $S_1 = 0$.

TERM OF DEGREE 2. Finally, $S_2 = C.Xx.Yy.Zz$, where

$$\begin{aligned} C &= [Xx, Yy][Xx, Zz] + [Xx, Yy][Yy, Zz] \\ &\quad + [Yy, Zz][Yy, Xx] + [Yy, Zz][Zz, Xx] \\ &\quad + [Zz, Xx][Zz, Yy] + [Zz, Xx][Xx, Yy] . \end{aligned}$$

Then $C = 0$ by the antisymmetry of the linking number. Thus $S_2 = 0$.

This concludes the proof of the Jacobi identity. Indeed

$$\begin{aligned} \{\{Xx, Yy\}_\alpha, Zz\}_\alpha + \{\{Yy, Zz\}_\alpha, Xx\}_\alpha + \{\{Zz, Xx\}_\alpha, Yy\}_\alpha \\ = \alpha^2 S_2 + \alpha S_1 + S_0 \\ = 0. \end{aligned}$$

□

2.3. The multi fraction algebra. The swapping algebra is very easy to define. However, in the sequel we shall need to consider other Poisson algebras built out of the swapping algebra and that will be more precisely subalgebras of the fraction algebra of $\mathcal{L}(\mathbf{P})$. We introduce in this paragraph *cross fractions*, *multi fractions* and the *multi fraction algebra*.

2.3.1. Cross fractions and multi fractions. Since $\mathcal{L}_\alpha(\mathbf{P})$ is an integral domain (with respect to the commutative product) we can consider its algebra of fractions $\mathcal{Q}_\alpha(\mathbf{P})$.

Let $(X, Y, x, y) =: Q$ be a quadruple of points of \mathbf{P} so that $x \neq Y$ and $y \neq X$. The *cross fraction* determined by Q is the element of $\mathcal{Q}_\alpha(\mathbf{P})$ defined by

$$[X; Y; x; y] := \frac{Xx.Yy}{Xy.Yx}.$$

More generally, let $\mathbf{X} := (X_1, \dots, X_n)$ and $\mathbf{x} := (x_1, \dots, x_n)$ are two tuples of elements of \mathbf{P} so that $x_i \neq X_i$ for all i , let σ be a permutation of $\{1, \dots, n\}$ then the *elementary multi fraction – defined over \mathbf{P}* – defined by this data is

$$[\mathbf{X}, \mathbf{x}; \sigma] := \frac{\prod_{i=1}^n X_i x_{\sigma(i)}}{\prod_{i=1}^n X_i x_i}.$$

2.3.2. The multi fraction algebra. Let now $\mathcal{B}(\mathbf{P})$ be the vector space generated by elementary multi fractions and let us call any element of $\mathcal{B}(\mathbf{P})$ a *multi fraction*. We have the following proposition

Proposition 2.3.1. *The vector space $\mathcal{B}(\mathbf{P})$ is a Poisson subalgebra of $\mathcal{Q}_\alpha(\mathbf{P})$. Moreover it is generated as a Poisson algebra by cross fractions. Finally the swapping bracket $\{\cdot, \cdot\}_\alpha$ when restricted to $\mathcal{B}(\mathbf{P})$ does not depend on α .*

From now on, we call the Poisson algebra $\mathcal{B}(\mathbf{P})$ the *algebra of multi fractions*.

Proof. The proposition follows from the following immediate observations:

- every elementary multi fraction is a product of cross fractions,
- if A and B are two cross fractions then $\{A, B\}_\alpha$ is a multi fraction and does not depend on α .

□

3. CROSS RATIOS AND CROSS FRACTIONS

In this section, we interpret cross fractions, and in general multi fractions, as functions on the space of cross ratios.

3.1. Cross ratios. Recall that a cross ratio on a set \mathbf{P} is a map \mathbf{b} from

$$\mathbf{P}^{4*} := \{(X, Y, x, y) \in \mathbf{P} \mid y \neq X, x \neq Y\}$$

to a field \mathbb{K} which satisfies some algebraic rules. Roughly speaking, these rules encode two conditions which constitute a normalisation, and two multiplicative cocycle identities which hold for some of the variables:

$$\text{NORMALISATION: } \mathbf{b}(X, Y, x, y) = 0 \Leftrightarrow x = X, \text{ or } Y = y,$$

$$\text{NORMALISATION: } \mathbf{b}(X, Y, x, y) = 1 \Leftrightarrow x = y, \text{ or } y = Y,$$

$$\text{COCYCLE IDENTITY: } \mathbf{b}(X, Y, x, y) = \mathbf{b}(X, Y, x, z)\mathbf{b}(X, Y, z, y),$$

$$\text{COCYCLE IDENTITY: } \mathbf{b}(X, Y, x, y) = \mathbf{b}(X, Z, x, y)\mathbf{b}(Z, W, x, y).$$

REMARKS:

- We change our convention from our previous articles [13, 14] in order to be coherent with the projective cross ratio: let \mathbf{b} be a cross ratio with respect to the definition above, let $b(X, x, Y, y) := \mathbf{b}(X, Y, x, y)$, then b is a cross ratio using our older convention.
- In the case \mathbf{P} has a Hölder structure and $\mathbb{K} = \mathbb{R}$, we assume furthermore assume that \mathbf{b} is Hölder.
- Assume Γ acts on \mathbf{P} . Let \mathbf{b} be a Γ invariant cross ratio. Let γ^+ and γ^- be two points of \mathbf{P} fixed by an element γ of Γ , then the following quantity

$$\mathbf{b}(\gamma^-, \gamma y, \gamma^+, y)$$

does not depends on the choice of y . In particular let S be a closed connected oriented surface of genus greater than 2, let \mathbf{P} be $\partial_\infty \pi_1(S)$ equipped with the action of $\pi_1(S)$. Let γ^+ and γ^- be respectively the attractive and repulsive fixed point of a non trivial element γ of Γ and \mathbf{b} a $\pi_1(S)$ -invariant cross ratio, then

$$\ell_{\mathbf{b}}(\gamma) := \left| \log \left| \mathbf{b}(\gamma^-, \gamma y, \gamma^+, y) \right| \right|,$$

is called the *period* of γ .

We finally denote by $\mathbb{B}(\mathbf{P})$ the set of cross ratios on \mathbf{P} .

These definitions are closely related to those given by Otal in [20, 21], those discussed from various perspectives by Ledrappier in [17] and those of Bourdon in [2] in the context of $\text{CAT}(-1)$ -spaces.

3.2. Multi fractions as functions. To every cross fraction $[X; Y; x; y]$, we associate a function, denoted by $\overline{[X; Y; x; y]}$, on $\mathbb{B}(\mathbf{P})$ by the following formula

$$\overline{[X; Y; x; y]}(\mathbf{b}) = \mathbf{b}(X, Y, x, y).$$

The following proposition follows at once from the definition of cross ratio

Proposition 3.2.1. *The map $[X; Y; x; y] \rightarrow \overline{[X; Y; x; y]}$ extends uniquely to a morphism of commutative associative algebras from $\mathcal{B}(\mathbf{P})$ to the algebra of functions on $\mathbb{B}(\mathbf{P})$.*

In the sequel, we shall use an identical notation for a multi fraction and its image in the space of functions on $\mathbb{B}(\mathbf{P})$. So far, we did not consider any topological structure on $\mathbb{B}(\mathbf{P})$.

3.3. Multi fractions and Hitchin components. In [13], we identified the Hitchin component with a space of cross ratios satisfying certain identities. Let us recall notations and definitions

3.3.1. Hitchin component. Let S be a closed oriented connected surface with genus at least two.

Definition 3.3.1. [FUCHSIAN AND HITCHIN HOMOMORPHISMS] *An n -Fuchsian homomorphism from $\pi_1(S)$ to $\text{PSL}_n(\mathbb{R})$ is a homomorphism ρ which factorises as $\rho = \iota \circ \rho_0$, where ρ_0 is a discrete faithful homomorphism with values in $\text{PSL}_2(\mathbb{R})$ and ι is an irreducible homomorphism from $\text{PSL}_2(\mathbb{R})$ to $\text{PSL}_n(\mathbb{R})$.*

An n -Hitchin homomorphism from $\pi_1(S)$ to $\text{PSL}_n(\mathbb{R})$ is a homomorphism which may be deformed into an n -Fuchsian homomorphism

The *Hitchin component* $\mathbf{H}(n, S)$ is the space of Hitchin homomorphisms up to conjugacy by an exterior automorphism of $\text{PSL}_n(\mathbb{R})$. By construction $\mathbf{H}(n, S)$ is identified with a connected component of the character variety. It is a result by Hitchin [12] that $\mathbf{H}(n, S)$ is homeomorphic to the interior of a ball of dimension $(3g - 2)n$.

As a corollary of the main result of [13], we have

Theorem 3.3.2. *If ρ is Hitchin, if γ is a non trivial element of $\pi_1(S)$ then $\rho(\gamma)$ has n distinct real eigenvalues of the same sign.*

This allows us to define the *width* of a non trivial element γ of $\pi_1(S)$ with respect to a Hitchin representation ρ as

$$\text{width}_\rho(\gamma) := \log \left(\left| \frac{\lambda_{\max}(\rho(\gamma))}{\lambda_{\min}(\rho(\gamma))} \right| \right),$$

where $\lambda_{\max}(\rho(\gamma))$ and $\lambda_{\min}(\rho(\gamma))$ are the eigenvalues of respectively maximum and minimum absolute values of the element $\rho(\gamma)$.

3.3.2. Rank n cross ratios. For every integer p , let $\partial_\infty \pi_1(S)_*^p$ be the set of pairs

$$(X, x) = ((X_0, X_1, \dots, X_p), (x_0, x_1, \dots, x_p)),$$

of $(p+1)$ -tuples of points in $\partial_\infty \pi_1(S)$ such that $X_j \neq X_i \neq x_0$ and $x_j \neq x_i \neq X_0$, whenever $j > i > 0$. Let $\chi^n(X, x)$ be the multi fraction defined by

$$\chi(X, x) := \det_{i,j>0} ([X_i; X_0; x_j; x_0]).$$

A cross ratio \mathbf{b} has *rank n* if

- $\chi^n(X, x)(\mathbf{b}) \neq 0$, for all (X, x) in $\partial_\infty \pi_1(S)_*^n$,
- $\chi^{n+1}(X, x)(\mathbf{b}) = 0$, for all (X, x) in $\partial_\infty \pi_1(S)_*^{n+1}$.

The main result of [15] – which used a result by Guichard [11] – is the following.

Theorem 3.3.3. *There exists a bijection ϕ from the set of n -Hitchin representations to the set of $\pi_1(S)$ -invariant rank n cross ratios, such that if $\mathbf{b} = \phi(\rho)$ then*

- (1) *for any nontrivial element γ of $\pi_1(S)$*

$$\ell_{\mathbf{b}}(\gamma) = \text{width}_\rho(\gamma),$$

where $\ell_{\mathbf{b}}(\gamma)$ is the period of γ is given with respect to $\mathbf{b} = \phi(\rho)$, and $\text{width}_\rho(\gamma)$ is the width of γ with respect to ρ .

- (2) *Moreover, if γ_1 and γ_2 are two non trivial elements of $\pi_1(S)$, if e_i , (respectively E_i) is an eigenvector of $\rho(\gamma_i)$ of maximal eigenvalue (respectively eigenvector of $\rho^*(\gamma_i)$ of minimum eigenvalue) then*

$$\mathbf{b}(\gamma_1^+, \gamma_2^+, \gamma_2^-, \gamma_1^-) = \frac{\langle E_2, e_1 \rangle \langle E_1, e_2 \rangle}{\langle E_1, e_1 \rangle \langle E_2, e_2 \rangle} \quad (30)$$

4. WILSON LOOPS, MULTI FRACTIONS AND LENGTH FUNCTIONS

In this section, we shall relate Wilson loops – which are regular functions on the character variety – to multi fractions. We in particular introduce *elementary functions* which are limits of Wilson loops, prove that they generate the multi fraction algebra and that they are smooth functions on the Hitchin component.

We also introduce length functions in Paragraph 4.4.

4.1. Wilson loops. Let γ be an element of $\pi_1(S)$ and ρ be an element of $\mathbf{H}(n, S)$. The *Wilson loop* associated to γ is the function $W(\gamma)$ on $\mathbf{H}(n, S)$ defined by

$$W(\gamma)(\rho) = \text{tr}(\rho(\gamma)).$$

We leave the reader check that this is indeed a well defined function.

Let us introduce the following definition.

Definition 4.1.1. [CLASS OF AN ELEMENT] *Let γ is a non trivial element of Γ , the class $[\gamma]$ of γ is the oriented pair (γ^+, γ^-) .*

Recall that $[\gamma] = [\eta]$ if and only if there exist positive integers m and n so that $\gamma^m = \eta^n$.

4.1.1. Asymptotics of Wilson loops. Let ρ be a Hitchin representation. Recall that for any γ in $\pi_1(S)$ we can write

$$\rho(\gamma) = \sum_{1 \leq i \leq n} \lambda_i(\gamma) p_i(\gamma),$$

where $p_i(\gamma)$ is a projector of trace 1, and $\lambda_i(\gamma)$ are real numbers so that

$$0 < |\lambda_n(\gamma)| < \dots < |\lambda_1(\gamma)|.$$

Let us denote $\dot{\rho}(\gamma) = p_1(\gamma)$

Definition 4.1.2. [GIRTH AND WIDTH] *Let the girth of ρ be*

$$\text{gh}(\rho) := \sup \left\{ \left| \frac{\lambda_k(\gamma)}{\lambda_j(\gamma)} \right| \mid k > j, \quad \gamma \in \rho(\pi_1(S)) \setminus \{\text{Id}\} \right\}, \quad (31)$$

and the width to be

$$\text{wh}(\rho) := \sup \left\{ \left| \frac{\lambda_j(\gamma)}{\lambda_k(\gamma)} \right| \mid \gamma \in \rho(\pi_1(S)) \setminus \{\text{Id}\} \right\}, \quad (32)$$

We have the following proposition

Proposition 4.1.3. *Let C be a compact subset of $\mathbf{H}(n, S)$ then*

$$\sup\{\text{gh}(\rho) \mid \rho \in C\} < 1.$$

Moreover there exists a constant W , we have

$$\sup\{\text{wh}(\rho) \mid \rho \in C\} < W.$$

Proof. This follows from the Anosov property. See also Sambarino ([?]) for details. \square

We choose an auxiliary norm, denoted $\|\cdot\|$ on \mathbb{R}^n . We define $[A]$ to be the set of eigenvectors of a purely loxodromic matrix A , and observe that $[A^n] = [A]$. Then we have

Proposition 4.1.4. *For any γ in $\rho(\pi_1(S))$, and $p \in \mathbb{N}$ we have*

$$\left\| \frac{\rho(\gamma^p)}{\mathbf{W}(\gamma^p)(\rho)} - \dot{\rho}(\gamma) \right\| \leq \text{gh}(\rho)^p K([\rho(\gamma)]), \quad (33)$$

where $K([\rho(\gamma)])$ only depends on $[\rho(\gamma)]$.

Proof. We can indeed write any real diagonalisable matrix with positive eigenvalues A as

$$A = \sum_{i=1}^{i=n} \lambda_i \mathbf{p}_i,$$

where \mathbf{p}_i are projectors. Thus

$$\begin{aligned} \left\| \frac{A^p}{\text{tr}(A^p)} - \mathbf{p}_1 \right\| &\leq \frac{1}{\sum_{i=1}^{i=n} \lambda_i^p} \left\| \sum_{i=2}^{i=n} \lambda_i^p \mathbf{p}_i - \left(\sum_{i=2}^{i=n} \lambda_i^p \right) \mathbf{p}_1 \right\| \\ &\leq \left(\frac{\lambda_2}{\lambda_1} \right)^p \left(n \|\mathbf{p}_1\| + \sum_{i=2}^{i=n} \|\mathbf{p}_2\| \right). \end{aligned} \quad (34)$$

Thus the inequality follows by taking

$$K([A]) = n \|\mathbf{p}_1\| + \sum_{i=2}^{i=n} \|\mathbf{p}_2\|.$$

\square

We begin with the following proposition.

Proposition 4.1.5. *Let $\gamma_1, \dots, \gamma_k$ be non trivial elements of $\pi_1(S)$. Then the sequence*

$$\left\{ \frac{\mathbf{W}(\gamma_1^p \dots \gamma_k^p)}{\mathbf{W}(\gamma_1^p) \dots \mathbf{W}(\gamma_k^p)} \right\}_{p \in \mathbb{N}}$$

converges uniformly on every compact of $\mathbf{H}(n, S)$ when p goes to infinity to a multi fraction. More precisely,

$$\lim_{p \rightarrow \infty} \left(\frac{\mathbf{W}(\gamma_1^p \dots \gamma_k^p)}{\mathbf{W}(\gamma_1^p) \dots \mathbf{W}(\gamma_k^p)} \right) = \frac{\prod_{i=1}^{i=k} \gamma_{i+1}^+ \gamma_i^-}{\prod_{i=1}^{i=k} \gamma_i^+ \gamma_i^-} = [\mathbf{G}^+, \mathbf{G}^-; \tau],$$

where $\mathbf{G}^\pm = (\gamma_1^\pm, \dots, \gamma_k^\pm)$ and $\tau(i) = i - 1$, using the convention that $k + 1 = 1$.

Proof. We first observe that if e_i , (respectively E_i) is an eigenvector of $\rho(\gamma_i)$ of maximal eigenvalue (respectively eigenvector of $\rho^*(\gamma_i)$ of minimum eigenvalue) so that $\langle E_i, e_i \rangle = 1$ then

$$\mathrm{tr}(\dot{\rho}(\gamma_1) \dots \dot{\rho}(\gamma_k)) = \prod_i \langle E_i, e_{i+1} \rangle.$$

By Equation (30),

$$\prod_i \langle E_i, e_{i+1} \rangle = \left(\frac{\prod_{i=1}^{i=p} \gamma_{i+1}^+ \gamma_i^-}{\prod_{i=1}^{i=k} \gamma_i^+ \gamma_i^-} \right) (\rho).$$

It thus follows that

$$\mathrm{tr}(\dot{\rho}(\gamma_1) \dots \dot{\rho}(\gamma_k)) = [\mathbf{G}^+, \mathbf{G}^-; \tau].$$

Then the result follows at once from Propositions 4.1.4 and 4.1.3. \square

4.2. Elementary functions. Proposition 4.1.5 leads us to the following definition.

Definition 4.2.1. *The multi fraction*

$$\mathbf{T}(\gamma_1, \dots, \gamma_p) := \frac{\prod_{i=1}^{i=k} \gamma_{i+1}^+ \gamma_i^-}{\prod_{i=1}^{i=k} \gamma_i^+ \gamma_i^-} \quad (35)$$

is an elementary function of order p

By the previous proposition and its proof, we have the following equalities

$$\mathbf{T}(\gamma_1, \dots, \gamma_p) = \lim_{n \rightarrow \infty} \frac{\mathbf{W}(\gamma_1^n \dots \gamma_p^n)}{\mathbf{W}(\gamma_1^n) \dots \mathbf{W}(\gamma_p^n)} \quad (36)$$

$$\mathbf{T}(\gamma_1, \dots, \gamma_p) = \mathrm{tr}(\dot{\rho}(\gamma_1) \dots \dot{\rho}(\gamma_p)). \quad (37)$$

The following formal properties of elementary functions are then easily checked.

Proposition 4.2.2. *The following properties of elementary functions hold*

- (1) CYCLIC INVARIANCE: *for every cyclic permutation σ of $\{1, \dots, p\}$ we have*

$$\mathsf{T}(\gamma_1, \dots, \gamma_p) = \mathsf{T}(\gamma_{\sigma(1)}, \dots, \gamma_{\sigma(p)})$$

- (2) CLASS INVARIANCE: *if $[\eta_i] = [\gamma_i]$ then*

$$\mathsf{T}(\gamma_1, \dots, \gamma_p) = \mathsf{T}(\eta_1, \dots, \eta_p).$$

- (3) *if $[\gamma_p] = [\gamma_{p-1}]$ then*

$$\mathsf{T}(\gamma_1, \dots, \gamma_p) = \mathsf{T}(\gamma_1, \dots, \gamma_{p-1})$$

- (4) *if $[\gamma_p] = [\gamma_{p-1}^{-1}]$ then*

$$\mathsf{T}(\gamma_1, \dots, \gamma_p) = 0$$

- (5) RELATIONS *Assume that $[\gamma_i] \neq [\gamma_{i+1}]$ then*

$$\mathsf{T}(\gamma_1, \dots, \gamma_p) = \frac{\mathsf{T}(\gamma_1, \gamma_2) \mathsf{T}(\gamma_1, \gamma_p) \mathsf{T}(\gamma_2, \gamma_3, \dots, \gamma_p)}{\mathsf{T}(\gamma_p, \gamma_2, \gamma_1)}$$

We deduce from the last statement the following corollary

Corollary 4.2.3. *Let P be the set of fixed points in $\partial_\infty \pi_1(S)$ of non trivial elements of $\pi_1(S)$. Then every restriction of an elementary multi fraction over P is a quotient of product of elementary functions of order 2 and 3.*

Proof. Let us consider a, b, c, d be four non trivial elements of $\pi_1(S)$, then we have

$$\frac{\mathsf{T}(a, b, c) \cdot \mathsf{T}(c, d)}{\mathsf{T}(a, d, c) \mathsf{T}(c, b)} = [b^+; d^+; a^-; c^-]. \quad (38)$$

The result follows. \square

Recall that in this section we choose P to be the set of fixed points of non trivial elements of $\pi_1(S)$. We now prove,

Proposition 4.2.4. *Every multi fraction – defined over P – is a smooth function on $\mathsf{H}(n, S)$.*

Proof. Let $\text{Hom}(n, S)$ be the space of Hitchin homomorphisms. Let π be the submersion

$$\pi : \text{Hom}(n, S) \rightarrow \mathsf{H}(n, S) = \text{Hom}(n, S) / \text{Aut}(\text{PSL}_n(\mathbb{R})).$$

For every loxodromic element A in $\text{PSL}_n(\mathbb{R})$ let p_A be the projection on the eigenspace of maximal eigenvalue with respect to the other

eigenspaces. The map $A \rightarrow p_A$ is smooth. It follows that for any elements $\gamma_1, \dots, \gamma_k$ in $\pi_1(S)$ the map from $\text{Hom}(n, S)$ to \mathbb{R} defined by

$$\Psi : \rho \rightarrow \text{tr}(p_{\rho(\gamma_1)} \cdots p_{\rho(\gamma_k)})$$

is smooth. We conclude by observing that Ψ is $\text{Aut}(\text{PSL}_n(\mathbb{R}))$ -invariant and that by Equation (37)

$$\Psi = \mathbb{T}(\gamma_1, \dots, \gamma_k) \circ \pi.$$

Thus every elementary function is smooth and by the previous result every multi fraction is smooth. \square

4.3. The swapping bracket of elementary functions. For the sequel, we shall need to compute the swapping brackets of elementary functions. This is given by the following proposition whose proof follows by an immediate application of the definition. We first say that two nontrivial elements γ and η in $\pi_1(S)$ are *coprime* if for all non zero integers m and n , $\gamma^n \neq \eta^m$.

Proposition 4.3.1. *Let $\gamma_0, \dots, \gamma_p$, respectively η_0, \dots, η_q be elements of $\pi_1(S) \setminus \{1\}$ such that (γ_i, γ_{i+1}) as well as (η_j, η_{j+1}) are pairwise coprime. Let*

$$\begin{aligned} a_{i,j} &:= [\gamma_i^+ \gamma_i^-, \eta_j^+ \eta_j^-] \\ b_{i,j} &:= [\gamma_{i+1}^+ \gamma_i^-, \eta_{j+1}^+ \eta_j^-] \\ c_{i,j} &:= [\gamma_i^+ \gamma_i^-, \eta_{j+1}^+ \eta_j^-] \\ d_{i,j} &:= [\gamma_{i+1}^+ \gamma_i^-, \eta_j^+ \eta_j^-] \\ \mathbb{T}_\gamma &:= \mathbb{T}(\gamma_0, \dots, \gamma_p) \\ \mathbb{T}_\eta &:= \mathbb{T}(\eta_0, \dots, \eta_q) \end{aligned} \tag{39}$$

Then

$$\begin{aligned} \frac{\{\mathbb{T}_\gamma, \mathbb{T}_\eta\}}{\mathbb{T}_\gamma \cdot \mathbb{T}_\eta} &= \sum_{\substack{0 \leq i \leq p, \\ 0 \leq j \leq q'}} \left(a_{i,j} \mathbb{T}(\gamma_i, \eta_j) + b_{i,j} \frac{\mathbb{T}(\eta_{j+1}, \eta_j, \gamma_{i+1}, \gamma_i)}{\mathbb{T}(\eta_j, \eta_{j+1}) \mathbb{T}(\gamma_i, \gamma_{i+1})} \right) \\ &- \sum_{\substack{0 \leq i \leq p, \\ 0 \leq j \leq q'}} \left(c_{i,j} \frac{\mathbb{T}(\gamma_i, \eta_{j+1}, \eta_j)}{\mathbb{T}(\eta_j, \eta_{j+1})} + d_{i,j} \frac{\mathbb{T}(\eta_j, \gamma_{i+1}, \gamma_i)}{\mathbb{T}(\gamma_i, \gamma_{i+1})} \right). \end{aligned} \tag{40}$$

Proof. Using “logarithmic derivatives”, we have

$$\begin{aligned}
\frac{\{\mathbb{T}_\gamma, \mathbb{T}_\eta\}}{\mathbb{T}_\gamma \cdot \mathbb{T}_\eta} &= \sum_{\substack{0 \leq i \leq p, \\ 0 \leq j \leq q}} \left(\left(\frac{\{\gamma_{i+1}^+ \gamma_i^-, \eta_{j+1}^+ \eta_j^-\}}{\gamma_{i+1}^+ \gamma_i^- \cdot \eta_{j+1}^+ \eta_j^-} + \frac{\{\gamma_i^+ \gamma_i^-, \eta_j^+ \eta_j^-\}}{\gamma_i^+ \gamma_i^- \cdot \eta_j^+ \eta_j^-} \right) \right. \\
&\quad \left. - \left(\frac{\{\gamma_i^+ \gamma_i^-, \eta_{j+1}^+ \eta_j^-\}}{\gamma_i^+ \gamma_i^- \cdot \eta_{j+1}^+ \eta_j^-} + \frac{\{\gamma_{i+1}^+ \gamma_i^-, \eta_j^+ \eta_j^-\}}{\gamma_{i+1}^+ \gamma_i^- \cdot \eta_j^+ \eta_j^-} \right) \right) \\
&= \sum_{\substack{0 \leq i \leq q \\ 0 \leq j \leq q'}} \left(b_{i,j} \frac{\gamma_{i+1}^+ \eta_j^- \cdot \eta_{j+1}^+ \gamma_i^-}{\gamma_{i+1}^+ \gamma_i^- \cdot \eta_{j+1}^+ \eta_j^-} + a_{i,j} \frac{\gamma_i^+ \eta_j^- \cdot \eta_j^+ \gamma_i^-}{\gamma_i^+ \gamma_i^- \cdot \eta_j^+ \eta_j^-} \right. \\
&\quad \left. - d_{i,j} \frac{\gamma_{i+1}^+ \eta_j^- \cdot \eta_j^+ \gamma_i^-}{\gamma_{i+1}^+ \gamma_i^- \cdot \eta_j^+ \eta_j^-} - c_{i,j} \frac{\gamma_i^+ \eta_j^- \cdot \eta_{j+1}^+ \gamma_i^-}{\gamma_i^+ \gamma_i^- \cdot \eta_{j+1}^+ \eta_j^-} \right).
\end{aligned}$$

From the definition of elementary functions (35), we get that

$$\begin{aligned}
\frac{\mathbb{T}(\eta_{j+1}, \eta_j, \gamma_{i+1}, \gamma_i)}{\mathbb{T}(\eta_j, \eta_{j+1}) \mathbb{T}(\gamma_i, \gamma_{i+1})} &= \frac{\gamma_{i+1}^+ \eta_j^- \cdot \eta_{j+1}^+ \gamma_i^-}{\gamma_{i+1}^+ \gamma_i^- \cdot \eta_{j+1}^+ \eta_j^-}, \\
\mathbb{T}(\gamma_i, \eta_j) &= \frac{\gamma_i^+ \eta_j^- \cdot \eta_j^+ \gamma_i^-}{\gamma_i^+ \gamma_i^- \cdot \eta_j^+ \eta_j^-}, \\
\frac{\mathbb{T}(\eta_j, \gamma_{i+1}, \gamma_i)}{\mathbb{T}(\gamma_i, \gamma_{i+1})} &= \frac{\gamma_{i+1}^+ \eta_j^- \cdot \eta_j^+ \gamma_i^-}{\gamma_{i+1}^+ \gamma_i^- \cdot \eta_j^+ \eta_j^-}, \\
\frac{\mathbb{T}(\gamma_i, \eta_{j+1}, \eta_j)}{\mathbb{T}(\eta_j, \eta_{j+1})} &= \frac{\gamma_i^+ \eta_j^- \cdot \eta_{j+1}^+ \gamma_i^-}{\gamma_i^+ \gamma_i^- \cdot \eta_{j+1}^+ \eta_j^-}.
\end{aligned}$$

This concludes the proof of the proposition \square

4.4. Length functions. We introduce in this paragraph length functions.

4.4.1. Length functions from the point of view of the multi fraction algebra. For any $y \in \partial_\infty \pi_1(S)$, let us introduce the following cross fraction,

$$p_\beta(y) = \frac{\beta^+ \beta^{-1}(y) \cdot \beta^- \beta(y)}{\beta^+ \beta(y) \cdot \beta^- \beta^{-1}(y)}.$$

We have, for any β in $\pi_1(S)$

$$p_\beta(y) - p_\beta(z) = F_{y,z} - (\beta^2)^* F_{y,z},$$

where

$$F_{y,z} = \frac{\beta^+ \beta^{-1} y \cdot \beta^- \beta^{-1}(z)}{\beta^+ \beta^{-1}(z) \cdot \beta^- \beta^{-1}(y)}$$

In particular, the restriction of $p_\beta(y)$ to the space of $\pi_1(S)$ -invariant cross ratios is independent on the choice of y .

For the sake of simplicity, we introduce the following formal series of multi fractions and call it a *length function*.

$$\hat{\ell}_\beta(y) := \log(p_\beta(y)),$$

extending the bracket by the “log derivative” formulas

$$\{\hat{\ell}_\beta(y), q\} := \frac{\{p_\beta(y), q\}}{p_\beta(y)}, \quad \{\hat{\ell}_\beta(y), \hat{\ell}_\gamma(z)\} := \frac{\{p_\beta(y), p_\gamma(z)\}}{p_\beta(y) \cdot p_\gamma(z)}. \quad (41)$$

Observe that $I_S(\hat{\ell}_{\beta^n}(y)) = n \cdot I_S(\hat{\ell}_\beta(y))$.

4.4.2. *Length functions and the character variety.* We can further relate these objects with the period and length defined in Paragraph 3.1.

Let

$$I_S : \mathcal{B}(\partial_\infty \pi_1(S)) \rightarrow C^\infty(\mathcal{H}(n, S)),$$

denote the restriction of functions from $\mathbb{B}(\partial_\infty \pi_1(S))$ to $\mathcal{H}(n)$.

We have that

$$\log I_S(\hat{\ell}_\beta(y)) = \ell_\beta,$$

where

$$\ell_\beta(\rho) = \ell_b(\beta),$$

where ℓ_b is the period of β which respect to the cross ratio associated to ρ (see Section 3.1).

5. THE GOLDMAN ALGEBRA

In this section, we first recall the construction of the Atiyah–Bott–Goldman symplectic form on the character variety. We then explain the construction of the Goldman algebra which allows to compute the bracket of Wilson loops in terms of a Lie bracket on the vector space generated by free homotopy classes of loops.

5.1. The Atiyah–Bott–Goldman symplectic form. In [1], Atiyah and Bott introduced a symplectic struture on the character variety of representations of closed surface groups in compact Lie group, generalizing Poincaré duality. This was later generalised by Goldman for non compact groups in [7, 6] and connected to the Weil–Petersson Kähler form. If identify the tangent space of $\mathcal{H}(n, S)$ at ρ with $H_\rho^1(\mathfrak{g})$, where \mathfrak{g} is the Lie algebra of $\mathrm{PSL}_n(\mathbb{R})$ then the symplectic form is given

$$\omega_S([A], [B]) = \int_V \mathrm{tr}(A \wedge B), \quad (42)$$

where A and B are de Rham representatives of the classes $[A]$ and $[B]$. We denote by $\{, \}_S$ the associated Poisson bracket, called the *Atiyah–Bott–Goldman (ABG) Poisson bracket* in the sequel, and $\mathcal{A}(S)$ the Poisson algebra of functions on $H(n, S)$. In the next paragraph, we show how to compute the Atiyah–Bott–Goldman bracket in the case of $\mathrm{PSL}_n(\mathbb{R})$ for the Wilson loops we introduced in the previous section.

5.2. Wilson loops and the Goldman algebra. We describe in this paragraph the Goldman algebra and how it helps computing the ABG-Poisson bracket. Let \mathcal{C} be the set of free homotopy class of closed curves on an oriented surface S . Let $\mathbb{Q}[\mathcal{C}]$ be the vector space generated by \mathcal{C} over \mathbb{Q} . We extend linearly Wilson loops so that the map $\gamma \mapsto w_\gamma$ is now a linear map from $\mathbb{Q}[\mathcal{C}]$ to $C^\infty(H(n, S))$.

Goldman introduced in [7] a Lie bracket on $\mathbb{Q}[\mathcal{C}]$. We define it for two elements γ_1 and γ_2 of $\mathcal{C} \subset \mathbb{Q}[\mathcal{C}]$ and then extend it to $\mathbb{Q}[\mathcal{C}]$ linearly. We choose two curves representing γ_1 and γ_2 , which we denote the same way.

If γ_1 and γ_2 are two curves from S^1 to S , an *intersection point* is a pair (a, b) in $S^1 \times S^1$ so that $\gamma_1(a) = \gamma_2(b)$. By a slight abuse of language, we usually identify an intersection point (a, b) with its image $x = \gamma_1(a) = \gamma_2(b)$. We further assume that γ_1 and γ_2 have transverse intersection points.

For every intersection point x , let ι_x be the local linking number at x , let $\gamma_1 *_x \gamma_2$ be the free homotopy class of the curve obtained by composing γ_1 and γ_2 in $\pi_1(S, x)$ and finally let

$$\iota(\gamma_1, \gamma_2) := \sum_{x \in \gamma_1 \cap \gamma_2} \iota_x,$$


be the global linking number.

Definition 5.2.1. *The Goldman bracket of γ_1 and γ_2 is the element of $\mathbb{Q}[\mathcal{C}]$ defined by*

$$\{\gamma_1, \gamma_2\} = \sum_{x \in \gamma_1 \cap \gamma_2} \iota_x \cdot \gamma_1 *_x \gamma_2. \quad (43)$$

We illustrate in Picture 2, the Goldman bracket of two curves. Goldman proved [7] that this bracket does not depend on the choice of representatives and is a Lie bracket. Moreover this bracket is related to the ABG-Poisson bracket as follows.

Theorem 5.2.2. [GOLDMAN] *Let γ_1 and γ_2 be two loops on S . Then the ABG-Poisson bracket of the two corresponding Wilson loops in*

FIGURE 2. $\{b, a\} = \alpha - \beta$

$H(n, S)$ is

$$\{W(\gamma_1), W(\gamma_2)\}_S = W(\{\gamma_1, \gamma_2\}) - \frac{\iota(\gamma_1, \gamma_2)}{n} W(\gamma_1) \cdot W(\gamma_2). \quad (44)$$

We just stated Goldman theorem for the case of $H(n, S)$, but the theorem has a formulation in the general case of character varieties for semi-simple groups.

6. VANISHING SEQUENCES AND THE MAIN RESULTS

In this section, we first recall the definition of the length functions on the character varieties, introduce the notion of a vanishing sequence of finite index subgroups of a surface group and state our main results relating the swapping algebra to the Goldman algebra. All these results will be proved in Section 9.

Let as usual

$$I_S : \mathcal{B}(\partial_\infty \pi_1(S)) \rightarrow C^\infty(H(n, S)),$$

denote the restriction of functions from $\mathbb{B}(\partial_\infty \pi_1(S))$ to $\mathcal{H}(n)$.

6.1. Poisson brackets of length functions. We explain in this section our results concerning length functions (See Paragraph 4.4 for the notation and definition). Our first result relates the Goldman and the swapping Poisson bracket.

Theorem 6.1.1. *Let γ and η be two geodesics with at most one intersection point, then we have*

$$\lim_{n \rightarrow \infty} I_S(\{\hat{\ell}_{\gamma^n}(y), \hat{\ell}_{\eta^n}(y)\}) = \{\ell_\gamma, \ell_\eta\}_S$$

In the course of the proof of this result, we prove the following result of independent interest which is an extension of Wolpert Formula [26, 25]

Theorem 6.1.2. [GENERALISED WOLPERT FORMULA].

Let γ and η two closed geodesics with a unique intersection point then the Goldman bracket of the two length functions ℓ_γ and ℓ_η seen as functions on the Hitchin component is

$$\{\ell_\gamma, \ell_\eta\}_S = \iota(\gamma, \eta) \sum_{\varepsilon, \varepsilon' \in \{-1, 1\}} \varepsilon \cdot \varepsilon' \cdot \mathsf{T}(\gamma^\varepsilon \cdot \eta^{\varepsilon'}), \quad (45)$$

where we recall that

$$\mathsf{T}(\xi, \zeta)(\rho) = \mathbf{b}_\rho(\xi^+, \zeta^+, \zeta^-, \xi^-).$$

We prove these two results in Paragraph 9.2

6.2. Poisson brackets of multi fractions. We now relate in general the swapping bracket and the Goldman bracket. Our result can be described by saying that the swapping bracket is an inverse limit (with respect to sequences of covering) of the Goldman bracket, or in other words that the swapping racket is a universal (in genus) Goldman bracket.

6.2.1. Vanishing sequences. Let S be a compact hyperbolic surface that is equipped with a hyperbolic metric. Let $\Gamma := \pi_1(S)$, let \tilde{S} be the universal cover of S so that $S = \pi_1(S)$. For any γ in $\pi_1(S)$, we denote by $\tilde{\gamma}$ its axis in \tilde{S} and $\langle \gamma \rangle$ the cyclic subgroup that it generates. Recall that we say that two elements γ and η of Γ are *coprime* if $\langle \gamma \rangle \cap \langle \eta \rangle = \{1\}$.

Let $\{\Gamma_m\}_{m \in \mathbb{N}}$ be a sequence of nested finite index subgroups of $\Gamma_0 := \Gamma$. Then let $S_n := \tilde{S}/\Gamma_n$. For any $\gamma \in \Gamma$ let $\langle \gamma \rangle_n := \langle \gamma \rangle \cap \Gamma_n$. Finally, let π_n be the projection from \tilde{S} to S_n and let $\tilde{\gamma}_n := \pi_n(\tilde{\gamma})$.

Definition 6.2.1. Let $\{\Gamma_m\}_{m \in \mathbb{N}}$ be a sequence of nested finite index normal subgroups of $\Gamma_0 := \Gamma$. We say that $\{\Gamma_m\}_{m \in \mathbb{N}}$ is a *vanishing sequence* if for all γ and η in Γ in Γ , for any set H invariant by left multiplication by γ and right multiplication by η whose projection in $\langle \eta \rangle \backslash \Gamma / \langle \gamma \rangle$ is finite, there exists n_0 , such that for all $n > n_0$, $H \subset \Gamma_n \cap \langle \eta \rangle \cdot \langle \gamma \rangle$.

We shall use freely the following immediate consequence

Proposition 6.2.2. For any η and γ in Γ , for any finite subset H_0 of Γ so that $H_0 \cap \langle \eta \rangle \cdot \langle \gamma \rangle = \emptyset$, there exists p_0 so that for all $p > p_0$, then

$$H \cap (\langle \eta \rangle \cdot \Gamma_p \cdot \langle \gamma \rangle) = \emptyset.$$

We prove in Appendix 11 that vanishing sequences exist. This is an immediate consequence of result by Niblo [19].

6.2.2. Sequences of subgroups and limits. Let \mathbf{P} be the subset of $\partial_\infty \pi_1(S)$ given by the end points of periodic geodesics. Let \mathbf{G} be the set of pairs of points $\gamma = (\gamma^-, \gamma^+)$ in \mathbf{P} which correspond to fixed points of by an element of the group $\partial_\infty(\pi_1(S))$. Observe that given any finite index subgroup Γ of $\pi_1(S)$, the set \mathbf{G} is in bijection with the set of primitive elements of Γ .

In the sequel, we shall freely identify elements of \mathbf{G} with primitive elements in any finite index subgroup of Γ

We associate to a sequence $\sigma = \{\Gamma_m\}_{m \in \mathbb{N}}$ of finite index subgroups of $\pi_1(S)$ the inverse limit S_σ of $\{S_m := \tilde{S}/\Gamma_m\}_{m \in \mathbb{N}}$, where \tilde{S} is the universal cover of S .

Observe that we have a map \mathbf{l} from $\mathcal{B}(\mathbf{P})$ to $\mathcal{A}(S_\sigma)$ which by definition is the projective limit of $\{\mathcal{A}(S_m)\}_{m \in \mathbb{N}}$.

Definition 6.2.3. Let $\{g_m\}_{m \in \mathbb{N}}$ be a sequence of functions, so that $g_m \in \mathcal{A}(S_m)$, we say that $\{g_m\}_{m \in \mathbb{N}}$ converges to the function h in $\mathcal{A}(S_\sigma)$ and write

$$\lim_{m \rightarrow \infty} g_m = h,$$

if for all p

$$\lim_{n \rightarrow \infty} \mathbf{l}_{S_p}(g_n) = \mathbf{l}_{S_p}(h),$$

where \mathbf{l}_{S_p} is the restriction with value in $\mathcal{A}(S_p)$.

6.2.3. Poisson brackets of multi fractions. The following result explains that the algebra of multi fractions is an inverse limit of Goldman algebras with respect to vanishing sequences .

Theorem 6.2.4. Let $\{\Gamma_m\}_{m \in \mathbb{N}}$ be a vanishing sequence of subgroups of $\pi_1(S)$. Let $\mathbf{P} \subset \partial_\infty \pi_1(S)$ be the set of end points of geodesics. Let b_0 and b_1 be two multi fractions in $\mathcal{B}(\mathbf{P})$. Then we have

$$\lim_{n \rightarrow \infty} \{\mathbf{l}(b_0), \mathbf{l}(b_1)\}_{S_n} = \mathbf{l}(\{b_0, b_1\}_W).$$

We prove this result in Paragraph 9.1.

7. PRODUCT FORMULAS AND BOUQUET IN GOOD POSITION

In this section, we wish to describe the Goldman bracket of curves which are compositions of many arcs. We shall call such a description a *product formula* and produce several instances of such formulas. This section is part of the technical core of this article.

The first formula – see Proposition 7.2.1 – deals with a rather general situation computing the Goldman bracket of curves which are compositions of many arcs. Then, considering repetition, and using special collection of arcs called *bouquets in good positions* – see Definition 7.3.2 – we prove a refinement of the product formula in Proposition 7.3.3. Proposition 7.3.3 is the first key result of this section.

Finally, in Proposition 7.5.2, we explain under which topological condition we can find bouquet in good position and compute the various intersection numbers involved in Proposition 7.3.3. Proposition 7.5.2 is the second key result of this section.

7.1. An alternative formulation of the Goldman bracket. We first need to give an alternative description of the Goldman bracket.

Let $\bar{\gamma}_1$ and $\bar{\gamma}_2$ be two arcs passing through a base point x_0 . For any point x in $\bar{\gamma}_i$, let $a_i(x)$ be the path along γ_i joining x_0 to x .

Definition 7.1.1. [INTERSECTION LOOPS] *Following these notations, for any $x \in \bar{\gamma}_1 \cap \bar{\gamma}_2$, the homotopy class*

$$c_x(\bar{\gamma}_1, \bar{\gamma}_2) := a_1(x).a_2(x)^{-1} \in \pi_1(S, x_0)$$

is called an intersection loop at x . – see Picture 3.




FIGURE 3. Intersection loop

The goal of this paragraph is the following proposition

Proposition 7.1.2. *Let γ_1 and γ_2 be two free homotopy classes of loops represented by curves $\bar{\gamma}_1$ and $\bar{\gamma}_2$ passing through x_0 . Then, the Goldman bracket in $\mathbb{Q}[\mathcal{C}]$ of the associated loops is given using intersection loops by*

$$\{\gamma_1, \gamma_2\}_S = \sum_{x \in \gamma_1 \cap \gamma_2} \iota(x) \bar{\gamma}_1 . c_x . \bar{\gamma}_2 . c_x^{-1}. \quad (46)$$

This proposition is an immediate consequence of the following

Proposition 7.1.3. *Let γ_1 and γ_2 be two loops passing through x_0 . Then for every $x \in \gamma_1 \cap \gamma_2$, we have*

$$\gamma_1 *_x \gamma_2 = \gamma_1 \cdot c_x(\gamma_1, \gamma_2) \gamma_2 \cdot c_x(\gamma_1, \gamma_2)^{-1},$$

as free homotopy classes of curves

Proof. Let as before a_i the arc along γ_i joining x_0 to x and $c_x = a_1 \cdot a_2^{-1}$, then

$$\begin{aligned} \gamma_1 *_x \gamma_2 &= a_1^{-1} \gamma_1 a_1 a_2^{-1} \gamma_2 a_2 \\ &= a_1^{-1} \gamma_1 c_x \gamma_2 a_2 \\ &= a_1^{-1} \gamma_1 c_x \gamma_2 c_x^{-1} a_1 \end{aligned}$$

Thus $\gamma_1 *_x \gamma_2$ is freely homotopic to $\gamma_1 c_x \gamma_2 c_x^{-1}$ □

7.2. The product formula. We need to express Goldman bracket of Wilson loop of curves consisting of many arcs. We work with the following data, see Figure 4 for a partial drawing,

- Let ξ_0, \dots, ξ_q and $\zeta_0, \dots, \zeta_{q'}$ be two tuples of arcs so that $A = \xi_0 \dots \xi_q$ and $B = \zeta_0 \dots \zeta_{q'}$ are closed curves.
- Assume furthermore that for all pairs (i, j) , ξ_i and ζ_j have transverse intersections and do not intersect at their end points.
- Let u_i – respectively v_i – be arcs joining a base point x_0 to the origin of ξ_i – respectively ζ_i .

Let us introduce the following notation


- for every $x \in \xi_i \cap \zeta_j$, let $c_x^{i,j} := c_x(u_i \xi_i, v_j \zeta_j)$,
- for any $\xi \in \pi_1(S)$, let $I_{i,j}(\xi) := \sum_{x \in \xi_i \cap \xi_j | \xi = c_x^{i,j}} \iota(x)$,
- let us denote $A_i := u_i \xi_i \xi_{i+1} \dots \xi_{i-1} u_i^{-1}$ and $B_j := v_j \zeta_j \zeta_{j+1} \dots \zeta_{j-1} v_j^{-1}$.

Proposition 7.2.1. [PRODUCT FORMULA] *Using the notation and assumptions described above, we have the following equality in $\mathbb{Q}[\mathcal{C}]$,*

$$\{A, B\} = \sum_{\substack{0 \leq i \leq q \\ 0 \leq j \leq q'}} \left(\sum_{x \in \xi_i \cap \zeta_j} \iota(x) A_i \cdot c_x^{i,j} \cdot B_j (c_x^{i,j})^{-1} \right) \quad (47)$$

$$= \sum_{\substack{0 \leq i \leq q \\ 0 \leq j \leq q'}} \left(\sum_{\xi \in \pi_1(S)} I_{i,j}(\xi) A_i \cdot \xi \cdot B_j \xi^{-1} \right) \quad (48)$$

We first prove a preliminary proposition and postpone the proof of Proposition 7.2.1 until the next paragraph.

FIGURE 4. Arcs ξ_i and u_i

7.2.1. *A preliminary case.* We first study the following simple situation

- Let ξ and η be two closed curves. Assume that $\xi = \xi_1 \cdot \xi_2$ and $\zeta = \zeta_1 \cdot \zeta_2$. Assume that of all i, j , ξ_i and ζ_j are closed curves with transverse intersections that do not intersect at their origin.
- Let u_i and v_j be arcs from x_0 to ξ_i and η_j respectively.
- Let $\tilde{\xi}_i := u_i \xi_i u_i^{-1}$, $\tilde{\zeta}_j := v_j \zeta_j v_j^{-1}$ and $c_x^{i,j} := c_x(\tilde{\xi}_i, \tilde{\zeta}_j) \in \pi_1(S, x_0)$ for $x \in \xi_i \cap \zeta_j$.

Proposition 7.2.2. *We have the following equality in $\mathbb{Q}[\mathcal{C}]$,*

$$\sum_{x \in \xi \cap \zeta} \iota(x) \xi *_{x} \zeta = \sum_{\substack{x \in \xi_i \cap \zeta_j \\ 1 \leq i, j \leq 2}} \iota(x) \tilde{\xi}_i \cdot \tilde{\xi}_{i+1} \cdot c_x^{i,j} \cdot \tilde{\zeta}_j \cdot \tilde{\zeta}_{j+1} \cdot (c_x^{i,j})^{-1}. \quad (49)$$

Proof. First, we observe that for any two pairs of curves (ξ_1, ξ_2) and (ζ_1, ζ_2) we have

$$(\xi_1 \cdot \xi_2) \cap (\zeta_1 \cdot \zeta_2) = \bigsqcup_{i,j} (\xi_i \cap \zeta_j).$$

Let us denote

$$c_x := c_x(\tilde{\xi}_1 \cdot \tilde{\xi}_2, \tilde{\zeta}_1 \cdot \tilde{\zeta}_2).$$

We then have

$$\begin{aligned} x \in \xi_1 \cap \zeta_1 &\implies c_x = c_x^{1,1}, \\ x \in \xi_2 \cap \zeta_1 &\implies c_x = \tilde{\xi}_1 \cdot c_x^{2,1}, \\ x \in \xi_1 \cap \zeta_2 &\implies c_x = c_x^{1,2} \cdot \tilde{\zeta}_1^{-1}, \\ x \in \xi_2 \cap \zeta_2 &\implies c_x = \tilde{\xi}_1 \cdot c_x^{2,2} \cdot \tilde{\zeta}_1^{-1}. \end{aligned}$$

Thus in all cases, if $x \in \xi_i \cap \zeta_j$ we have the following equality of free homotopy classes

$$\tilde{\xi}_1 \tilde{\xi}_2 c_x \tilde{\zeta}_1 \tilde{\zeta}_2 = \tilde{\xi}_i \tilde{\xi}_{i+1} \cdot c_x^{i,j} \cdot \tilde{\zeta}_j \cdot \tilde{\zeta}_{j+1} \cdot (c_x^{i,j})^{-1},$$

Thus we obtain the product formula.

$$\begin{aligned} \sum_{x \in (\xi_1 \xi_2) \cap (\zeta_1 \zeta_2)} \iota(x) \left(\tilde{\xi}_1 \cdot \tilde{\xi}_2 \cdot c_x \cdot \tilde{\zeta}_1 \cdot \tilde{\zeta}_2 \cdot c_x^{-1} \right) \\ = \sum_{i,j} \left(\sum_{x \in \xi_i \cap \zeta_j} \iota(x) (\tilde{\xi}_i \tilde{\xi}_{i+1} \cdot c_x^{i,j} \cdot \tilde{\zeta}_j \cdot \tilde{\zeta}_{j+1} \cdot (c_x^{i,j})^{-1}) \right). \end{aligned} \quad (50)$$

This concludes the proof \square

7.2.2. Proof of Proposition 7.2.1. Obviously Formula (48) is an immediate consequence of Formula (47), so we concentrate on the latter.

First, we observe that the product formula when ξ_i and ζ_j are closed curves follows by induction from Proposition 7.2.2.

Let us now make the following observation. Let a , ξ and ζ be three arcs, transverse to a curve κ . Assume that $\xi \cdot a \cdot a^{-1} \cdot \zeta$ is a closed curve, then we have the following equalities in $\mathbb{Q}[\mathcal{C}]$

$$\begin{aligned} \xi \cdot a \cdot a^{-1} \cdot \zeta &= \xi \cdot \zeta, \\ \sum_{x \in (\xi \cdot \zeta) \cap \kappa} \iota(x) \xi \cdot \zeta \cdot c_x \kappa \cdot c_x^{-1} &= \sum_{x \in (\xi \cdot a \cdot a^{-1} \cdot \zeta) \cap \kappa} \iota(x) \xi \cdot a \cdot a^{-1} \cdot \zeta \cdot c_x \kappa \cdot c_x^{-1}. \end{aligned} \quad (51)$$

The first equality is obvious. For the second we notice that every intersection point of a with κ appears twice with a different sign.

We can now extend the product formula to arcs: we choose auxiliary arcs α_i joining x_0 to the initial point of ξ_i , similarly auxiliary arcs β_i joining x_0 to the initial point of ζ_i and replace ξ_i by the closed curves $\hat{\xi}_i = \alpha_i \xi_i \alpha_{i+1}^{-1}$ and $\hat{\zeta}_i = \beta_i \zeta_i \beta_{i+1}^{-1}$ respectively. From Assertion (51), since the product formula holds for the closed curves $\hat{\zeta}_j$ and $\hat{\xi}_i$, it holds for the arcs ζ_j and ξ_i .

7.3. Bouquets in good position and the product formula. We shall need a special case of the product formula when we allow some repetitions in the arcs.

7.3.1. Bouquets in good position.

Definition 7.3.1. [FLOWERS AND BOUQUETS]

- (1) A flower based at (x_0, \dots, x_q) is a collection of arcs

$$\mathcal{S} := ((g_0, \dots, g_q), (\alpha_0, \dots, \alpha_q)),$$

such that

- g_i are closed curves based at x_i representing primitive elements in the fundamental group,
- α_i are arcs, called connecting arcs, joining x_i to x_{i+1} .

- (2) A bouquet is a triple

$$\mathcal{C} = (\mathcal{S}_0, \mathcal{S}_1, V),$$

where \mathcal{S}_1 and \mathcal{S}_0 are flowers based at (x_0, \dots, x_q) and $(y_0, \dots, y_{q'})$ respectively and V is an arc joining x_0 and y_0 .

- (3) We finally say that the bouquet \mathcal{C} represents $((\gamma_0, \dots, \gamma_q), (\eta_0, \dots, \eta_{q'}))$, where γ_i and η_j are the elements of $\pi_1(S, x_0)$ defined by $\gamma_i = U_i g_i U_i^{-1}$ and $\eta_j = V_j h_j V_j^{-1}$, where $U_i := \alpha_0 \dots \alpha_{i-1}$ and $V_j := V \cdot \beta_0 \dots \beta_{j-1}$,

We shall also need bouquets which have special configuration: let

$$\mathcal{C} = (((g_0, \dots, g_q), (\alpha_0, \dots, \alpha_q)), ((h_0, \dots, h_{q'}), (\beta_0, \dots, \beta_{q'})), V)$$

be a bouquet of flowers based respectively at (x_0, \dots, x_q) and $(y_0, \dots, y_{q'})$.

Definition 7.3.2. [GOOD POSITION] We say

- (1) \mathcal{C} is in a good position if
- the arcs α_i and g_i intersect transversally the arcs β_j and h_j at points different than x_i and y_j for all i, j ,
 - the closed curves $\alpha_0 \dots \alpha_q$ and $\beta_0 \dots \beta_{q'}$ are homotopic to zero.
- (2) \mathcal{C} is in a homotopically good position if it is in a good position and if the following intersection loops are homotopically trivial

$$\begin{aligned} c_x(U_i \cdot \alpha_i, V_j \cdot \beta_j), & \text{ for } x \in \alpha_i \cap \beta_j \\ c_x(U_i \cdot \alpha_i, V_j \cdot h_j), & \text{ for } x \in \alpha_i \cap h_j \\ c_x(U_i \cdot g_i, V_j \cdot \beta_j), & \text{ for } x \in g_i \cap \beta_j, \end{aligned} \tag{52}$$

where $U_i := \alpha_0 \dots \alpha_{i-1}$ and $V_j := V \cdot \beta_0 \dots \beta_{j-1}$.

In Figure 5, we have represented two flowers, one in blue, the other in red where the connecting arcs α_i and β_i are dotted. In this figure all intersection loops corresponding to the four yellow transverse intersection points are drawn in the orange contractible region. Thus the bouquet is in a homotopically good position.




FIGURE 5. Bouquet in good position

7.3.2. *Product formula for bouquets.* Let \mathcal{C} be a bouquet as above in good position. Let us consider the closed curves

$$\begin{aligned}\mathbf{F}_i^{(p,m)} &:= U_i \cdot g_i^n \cdot (\alpha_i \cdot g_{i+1}^p \alpha_{i+1} \cdots g_{i-1}^p \alpha_{i-1}) g_i^{p-n} U_i^{-1}, \\ \mathbf{G}_i^{(p,m)} &:= V_i \cdot h_i^n \cdot (\beta_i \cdot h_{i+1}^p \beta_{i+1} \cdots h_{i-1}^p \beta_{i-1}) h_i^{p-n} V_i^{-1}.\end{aligned}$$

To simplify notation, let us write $\mathbf{F}^{(p)} := \mathbf{F}_0^{(p,0)}$ and $\mathbf{G}^{(p)} := \mathbf{G}_0^{(p)}$.

Let us denote

$$\begin{aligned}H_{i,j} &:= \{c_x(U_i \cdot g_i, V_j \cdot h_j) \mid x \in g_i \cap h_j, c_x(U_i \cdot g_i, V_j \cdot h_j) \text{ is homotopically trivial} \}, \\ C_{i,j} &:= \{c_x(U_i \cdot g_i, V_j \cdot h_j) \mid x \in g_i \cap h_j, c_x(U_i \cdot g_i, V_j \cdot h_j) \text{ not homotopically trivial} \},\end{aligned}$$

Let finally

$$\begin{aligned}f_{i,j}(\mathcal{C}) &:= \sum_{\xi \in H_{i,j}} I_{i,j}(\xi), & m_{i,j}(\mathcal{C}) &:= \iota(g_i, \beta_j), \\ n_{i,j}(\mathcal{C}) &:= \iota(\alpha_i, h_j), & q_{i,j}(\mathcal{C}) &:= \iota(\alpha_i, \beta_j),\end{aligned} \quad (53)$$

where we recall that for any $\xi \in \pi_1(S)$, we denote

$$I_{i,j}(\xi) = \sum_{x \in g_i \cap h_j \mid c_x(U_i \cdot g_i, V_j \cdot h_j) = \xi} \iota(x).$$

We can rewrite the product formula.

Proposition 7.3.3. [PRODUCT FORMULA IN GOOD POSITION] *Assuming the bouquet \mathcal{C} is in a homotopically good position and using the above notation, we have the following equality in $\mathbb{Q}[\mathcal{C}]$,*

$$\begin{aligned} \{\mathbf{F}^{(p)}, \mathbf{G}^{(p)}\} &= \sum_{\substack{0 \leq i \leq q \\ 0 \leq j \leq q'}} \left(\sum_{\substack{1 \leq m' \leq p \\ 1 \leq m \leq p}} \mathbf{f}_{i,j}(\mathcal{C}) \left(\mathbf{F}_i^{(p,m')} \mathbf{G}_j^{(p,m)} \right) + \sum_{1 \leq m \leq p} \mathbf{m}_{i,j}(\mathcal{C}) \left(\mathbf{F}_i^{(p,m)} \mathbf{G}_j^{(p,0)} \right) \right. \\ &\quad \left. + \sum_{1 \leq m' \leq p} \mathbf{n}_{i,j}(\mathcal{C}) \left(\mathbf{F}_i^{(p,0)} \mathbf{G}_j^{(p,m')} \right) + \mathbf{q}_{i,j}(\mathcal{C}) \left(\mathbf{F}_i^{(p,0)} \mathbf{G}_j^{(p,0)} \right) \right) \\ &\quad + \sum_{\substack{0 \leq i \leq q \\ 0 \leq j \leq q'}} \sum_{\xi \in C_{i,j}} I_{i,j}(\xi) \left(\sum_{\substack{1 \leq m' \leq p \\ 1 \leq m \leq p}} \left(\mathbf{F}_i^{(p,m')} \xi \mathbf{G}_j^{(p,m)} \xi^{-1} \right) \right). \end{aligned} \quad (54)$$

Proof. This will be just another way to write the product formula. We consider the arcs ξ_i defined by

- $\xi_i := g_j$ if $i = j.(p+1) + n$ with $1 \leq n \leq p$,
- $\xi_i := \alpha_j$ if $i = j(p+1)$.

Similarly, we consider the arcs ζ_i

- $\zeta_i := h_j$ if $i = j.(p+1) + n$ with $1 \leq n \leq p$,
- $\zeta_i := \beta_j$ if $i = j.(p+1)$.

Let now finally consider the following arcs,

- $u_i := U_j = \alpha_0 \dots \alpha_j$, if $i = j.(p+1) + n$ with $1 \leq n \leq p$,
- $v_i := V_j = \beta_0 \dots \beta_j$, if $i = j.(p+1) + n$ with $1 \leq n \leq p$,

so that u_i , respectively v_i , goes from x_0 to x_j , respectively x_0 to y_j .

We now apply Formulae (47) and (54) for the arcs ξ_i, u_i, ζ_j, v_j . Observe that using the notation of Paragraph 7.2, we have

$$\mathbf{F}^{(p)} = A, \quad \mathbf{G}^{(p)} = B.$$

We now have to identify the term in the righthand sides of Formulae (47) and (54), and in particular understand the arcs $A_i, B_j, c_x^{i,j}$ that appears in the righthand side of Formula (54). By definition

$$A_i = u_i \xi_i \xi_{i+1} \dots \xi_{i-1} u_i^{-1}.$$

Thus if $i = j(p+1) + m$ with $0 \leq m \leq p$

$$A_i = \mathbf{F}_j^{(p,m)},$$

and by a similar argument

$$B_i = \mathbf{G}_j^{(p,m)}.$$

By definition if $x \in \xi_i \cap \zeta_j$,

$$c_x^{i,j} = c_x(u_i \xi_i, v_j \zeta_j).$$

We now observe that,

- (1) if $i = j.(p+1)$, then $u_i \xi_i = U_j.g_j$,
- (2) if $i = j.(p+1) + n$ with $1 \leq m \leq p$, then $u_i \xi_i = U_j.\alpha_j$,

and similary

- (1) if $i = j.(p+1)$, then $v_i \zeta_i = V_j.h_j$,
- (2) if $i = j.(p+1) + n$ with $1 \leq m \leq p$, then $v_i \zeta_i = V_j.\beta_j$,

Then the special Product Formula (54) is a consequence of the Product Formula (47): indeed, thanks to the “homotopically good position” hypothesis, many of the intersection loops $c_x^{i,j}$ are homotopically trivial. \square

7.4. Bouquets and covering. Let $\pi : S_1 \rightarrow S_0$ be a finite covering. Let

$$\mathcal{C} = (((g_0, \dots, g_q), (\alpha_0, \dots, \alpha_q)), ((h_0, \dots, h_{q'}), (\beta_0, \dots, \beta_q)), V)$$

be a bouquet of flowers in S_0 based respectively at (x_0, \dots, x_q) and $(y_0, \dots, y_{q'})$.

Let \hat{x}_0 be a lift of x_0 in S_1 .

Definition 7.4.1. *The bouquet of flowers in S_1*

$$\hat{\mathcal{C}} = (((\hat{g}_0, \dots, \hat{g}_q), (\hat{\alpha}_0, \dots, \hat{\alpha}_q)), ((\hat{h}_0, \dots, \hat{h}_{q'}), (\hat{\beta}_0, \dots, \hat{\beta}_q)), \hat{V})$$

is the lift of \mathcal{C} through \hat{x}_0 if,

- all arcs \hat{V} , $\hat{\alpha}_i$ and $\hat{\beta}_i$ are lifts of the arcs V , α_i and β_i .
- \hat{g}_0 is based at \hat{x}_0 .
- the closed curves \hat{g}_i and \hat{h}_j are the primitive lifts of the curves g_i and h_j , in other words the primitive curves which are lift of positive powers of the curves g_i and h_j .

Observe that the lift of a bouquet in homotopically good position is in homotopically good position.

7.5. Finding bouquets in good position. Let S be a closed hyperbolic surface and \tilde{S} its universal cover. Let $G = (\gamma_0, \dots, \gamma_q)$ and $F = (\eta_0, \dots, \eta_{q'})$ be two tuples of primitive elements of $\pi_1(S)$ such that for all i , (γ_i, γ_{i+1}) , are pairwise coprime as well as (η_i, η_{i+1}) , where the index i lives in $\mathbb{Z}/q\mathbb{Z}$ and $\mathbb{Z}/q'\mathbb{Z}$ respectively. Recall that we denote by $\tilde{\zeta}$ the axis of the element $\zeta \in \pi_1(S)$.

Definition 7.5.1. *We say G and F satisfy the Good Position Hypothesis if there exists a metric ball B in \tilde{S} such that*

- (1) *for all i and j so that γ_i and η_j are coprime,*

$$\tilde{\gamma}_i \cap \tilde{\eta}_j \subset B \quad (55)$$

- (2) *for all $\xi \in \pi_1(S) \setminus \{1\}$ we have*

$$B \cap \xi(B) = \emptyset. \quad (56)$$

- (3) *for all $\zeta \in \{\gamma_0, \dots, \gamma_q, \eta_0, \dots, \eta_{q'}\}$, for all $\xi \in \pi_1(S) \setminus \langle \zeta \rangle$ we have*

$$B \cap \xi(\tilde{\zeta}) = \emptyset. \quad (57)$$

- (4) *for all $\zeta \in \{\gamma_0, \dots, \gamma_q, \eta_0, \dots, \eta_{q'}\}$, for all $\xi \in \pi_1(S) \setminus \langle \zeta \rangle$ we have*

$$\tilde{\zeta} \cap \xi(\tilde{\zeta}) = \emptyset. \quad (58)$$

In other words, the closed geodesic corresponding to ζ is embedded.

Then, we have the following result

Proposition 7.5.2. *With the notation above, assume that G , F and $\pi_1(S)$ satisfy the Good Position Hypothesis, then there exist two bouquets \mathcal{C}_L and \mathcal{C}_R in S in a homotopically good position, both representing (G, F) such that furthermore*

$$\frac{1}{2}(\mathbf{f}_{i,j}(\mathcal{C}_L) + \mathbf{f}_{i,j}(\mathcal{C}_R)) = [\gamma_i^- \gamma_i^+, \eta_j^- \eta_j^+], \quad (59)$$

$$\frac{1}{2}(\mathbf{n}_{i,j}(\mathcal{C}_L) + \mathbf{n}_{i,j}(\mathcal{C}_R)) = [\gamma_i^- \gamma_{i+1}^-, \eta_j^- \eta_j^+], \quad (60)$$

$$\frac{1}{2}(\mathbf{m}_{i,j}(\mathcal{C}_L) + \mathbf{m}_{i,j}(\mathcal{C}_R)) = [\gamma_i^- \gamma_i^+, \eta_j^- \eta_{j+1}^-], \quad (61)$$

$$\frac{1}{2}(\mathbf{q}_{i,j}(\mathcal{C}_L) + \mathbf{q}_{i,j}(\mathcal{C}_R)) = [\gamma_i^- \gamma_{i+1}^-, \eta_j^- \eta_{j+1}^-]. \quad (62)$$

Proof. Let G and F be as above and B be a metric ball in \tilde{S} satisfying the assumptions (55), (56) and (57). We subdivide the proof in several steps. We denote by π the projection from \tilde{S} to S .

STEP 1: CONSTRUCTION OF THE BOUQUET IN GOOD POSITION

Let $\tilde{\gamma}_i$ be the axis of γ_i , let ε be some constant that we shall choose later to be very small and $\tilde{\eta}_j^\varepsilon$ be a curve (with constant geodesic curvature) at distance ε of the axis $\tilde{\eta}_j$ of η_j (Notice that we have too such curves, for the moment we choose arbitrarily one of them). We choose ε small enough so that Assertions (55) and (57) still hold when $\tilde{\eta}_j$ are replaced by $\tilde{\eta}_j^\varepsilon$.

For every i , let $x_i \in \tilde{\gamma}_i \cap B$ so that

$$\tilde{\gamma}_i \cap B \subset [x_i, \gamma_i^+[,$$

similarly, let $y_j \in \tilde{\eta}_j^\varepsilon$ so that

$$\tilde{\eta}_j^\varepsilon \cap B \subset [y_j, \eta_j^+[\varepsilon,$$

where $[a, b]_\varepsilon$ denote an arc joining a to b along a curve at a distance ε to a geodesic (see Figure 6).




FIGURE 6. Finding a bouquet in good position

We now consider geodesics arcs $\tilde{\alpha}_i$, $\tilde{\beta}_j$ and \tilde{V} in \tilde{S} joining respectively x_i to x_{i+1} , y_j to y_{j+1} and x_0 to y_0 . We furthermore choose B (and ε) so that all the arcs $\tilde{\alpha}_i$, $\tilde{\eta}_j^\varepsilon$, $\tilde{\gamma}_j$ and $\tilde{\beta}_j$ are transverse. In particular, if

$$\begin{aligned} \alpha_i &= \pi(\tilde{\alpha}_i), & \beta_i &= \pi(\tilde{\beta}_i), & V &= \pi(\tilde{V}), \\ \tilde{g}_i &= [x_i, \gamma(x_i)], & \tilde{h}_j &= [y_j, \eta_j(y_j)]_\varepsilon, \\ g_i &= \pi([x_i, \gamma(x_i)]), & h_j &= \pi([y_j, \eta_j(y_j)]_\varepsilon), \end{aligned} \quad (63)$$

then

$$\mathcal{C} = (((g_0, \dots, g_q), (\alpha_0, \dots, \alpha_q)), ((h_0, \dots, h_{q'}), (\beta_0, \dots, \beta_{q'})), V)$$

is in good position. Observe furthermore that \mathcal{C} represents (G, F) .

STEP 2: HOMOTOPICALLY GOOD POSITION

Let us now prove that \mathcal{C} is in a homotopically good position. Let as usual

$$U_i = \alpha_0 \dots \alpha_{i-1}, \quad V_j = V \cdot \beta_0 \dots \beta_{j-1}, \quad (64)$$

and

$$\tilde{U}_i = \tilde{\alpha}_0 \dots \tilde{\alpha}_{i-1}, \quad \tilde{V}_j = \tilde{V} \cdot \tilde{\beta}_0 \dots \tilde{\beta}_{j-1}. \quad (65)$$

Then \tilde{U}_i and \tilde{V}_j are the lifts of U_i and V_j respectively starting from x_0 and y_0 an ending respectively in x_i and y_j .

Observe that all the arcs $\tilde{\alpha}_k$, $\tilde{\beta}_l$ and \tilde{V} lie in B . and thus so do the paths \tilde{U}_i and \tilde{V}_j .

Let W_i be equal to α_i or g_i . Let \widehat{W}_j be equal to β_j or h_j . Let us fix from now $x \in W_i \cap \widehat{W}_j$.

Let us introduce some notation,

- Let a be the path along W_i from $\pi(x_i)$ to x , and \hat{a} be the path along \widehat{W}_j from $\pi(y_j)$ point to x .
- Let b and \hat{b} the lifts of a and \hat{a} in \tilde{S} starting respectively from x_i and x_j ,
- Let finally z and \hat{z} be the endpoints of b and \hat{b} and $\zeta \in \pi_1(S)$ so that $z = \zeta(\hat{z})$.

By construction ζ is conjugated to the intersection loop $c_x(U_i W_i, U_j \widehat{W}_j)$.

Let us now consider the various possibility about the position of z and \hat{z} .

- (1) $W_i = \alpha_i$, then $b \subset \tilde{\alpha}_i$ and thus z belongs to B .
- (2) $W_i = g_i$, then $z \in [x_i, \gamma_i(x_i)] \subset [x_i, \gamma_i^+]$,
- (3) $\widehat{W}_j = \beta_j$, then, symmetrically, $\hat{z} = \zeta(z)$ belongs to B .
- (4) $\widehat{W}_j = h_j$, then, symmetrically, $\hat{z} \in [y_j, \eta_j^\varepsilon(x_j)]_\varepsilon \subset [y_j, \eta_j^+]$ (where the intervals are subsets of $\tilde{\eta}_j^\varepsilon$).

Our goal is now to prove that $\zeta = 1$ unless, maybe, $W_i = g_i$ and $\widehat{W}_j = h_j$.

- (1) $W_i = \alpha_i$ and $\widehat{W}_j = \beta_j$ then by Assertions (1) and (3) above, both z and $\zeta(z)$ belong to B , and thus by Assertion (56), $\zeta = 1$.
- (2) $W_i = \alpha_i$ and $\widehat{W}_j = h_j$, then by Assertions (1) and (4), $\zeta(z) \in \tilde{\eta}_j^\varepsilon$ and $z \in B$. Thus $\zeta^{-1}(\tilde{\eta}_i^\varepsilon) \cap B \neq \emptyset$. Then by Hypothesis (57), $\zeta \in \langle \eta_j \rangle$. In particular $z \in B \cap \tilde{\eta}_j^\varepsilon$ and thus we have

$$z \in [y_j, \eta_j(y_j)]_\varepsilon, \quad (66)$$

Recall that from Assertion (4)

$$\zeta(z) = \hat{z} \in [y_j, \eta_j(y_j)]_\varepsilon. \quad (67)$$

Since η_j is primitive and $\zeta \in \langle \eta_j \rangle$, we obtain from Assertions (66) and (67) that $\zeta = 1$.

- (3) A symmetric argument proves that when $W_i = g_i$ and $\widehat{W}_j = \beta_j$, then $\zeta = 1$

This finishes the proof that \mathcal{C} is in a homotopically good position.

STEP 3: COMPUTATION OF THE INTERSECTION NUMBERS

Recall that for each (oriented) axis $\tilde{\eta}_j$ we had two choices of curves at distance ε . Let us denote by $\tilde{\eta}_j^L$ – respectively $\tilde{\eta}_j^R$ – the curve on the left –respectively on the right– to $\tilde{\eta}_j$. Let then \mathcal{C}^L and \mathcal{C}^R the corresponding collections of arcs.

We have proved that both \mathcal{C}^L and \mathcal{C}^R are in homotopically good position. Let us now compute the intersection numbers. We will do that step by step.

We shall repeat the following observation several time: let g and h be two curves in S passing through a point x_0 , intersecting transversally a finite number of points x_1, \dots, x_n . Let \tilde{g} and \tilde{h} the lift of these curves in \tilde{S} passing through a point \tilde{x}_0 . Then there the projection realizes a bijection between the set of theses x_i whose intersection loop is trivial, and the intersections points of \tilde{g} and \tilde{h} .

In particular

$$\sum_{x \in g \cap h \mid c_x(g, h) = 1} \iota(x) = \sum_{z \in \tilde{g} \cap \tilde{h}} \iota(z) \quad (68)$$

Proof of Equation (59): If γ_i and η_j are coprime, by Formula (68) and since two geodesics have at most one intersection point, we have that

$$f_{i,j} = [\gamma_i^- \gamma_i^+, \eta_j^- \eta_j^+].$$

If γ_i and η_j are not coprime, since g_i is embedded by Assumption (58), it follows that

$$\iota(g_i, h_j) = 0 = [\gamma_i^- \gamma_i^+, \eta_j^- \eta_j^+].$$

Thus in both cases

$$f_{i,j}(\mathcal{C}^L) = f_{i,j}(\mathcal{C}^R) = [\gamma_i^- \gamma_i^+, \eta_j^- \eta_j^+].$$

Proof of Equation (60): Since all the corresponding intersection loops are trivial, we see that

$$\iota(g_i, \beta_j) = \iota(\tilde{\gamma}_i, \tilde{\beta}_j).$$

We know that $\tilde{\beta}_j \subset B$. To simplify, let us first consider the case when γ_i and η_k are coprime for $k = j, j+1$. Then $\tilde{\gamma}_i \cap \tilde{\eta}_k^\varepsilon \subset B$ and thus

$$\iota(\tilde{\gamma}_i,]\eta_k^-, y_k[_\varepsilon) = 0.$$

It follows then that

$$\iota(g_i, \beta_j) = \iota(\tilde{\gamma}_i,]\eta_j^-, y_j[_\varepsilon \cup \tilde{\beta}_j \cup]y_{j+1}, \eta_{j+1}^-[_\varepsilon) = [\gamma_i^- \gamma_i^+, \eta_j^- \eta_{j+1}^-].$$

We illustrate that situation in Figure 7a




FIGURE 7. Intersection computations

Let us move to the remaining cases. The purpose of taking the “left and right perturbations” of $\tilde{\eta}_j$ is to take care of the situation when η_j (or η_{j+1}) and γ_i are not coprime. So let us assume now that $\tilde{\eta}_j = \tilde{\gamma}_i$ (the case when $\tilde{\eta}_{j+1} = \tilde{\gamma}_i$ is symmetric).

Then in this case assume that η_{j+1}^- is on the left of $\tilde{\eta}_j$ and $\tilde{\gamma}_i$ has the same orientation as $\tilde{\eta}_j$ (the other cases being symmetric). It then follows that

$$m_{i,j}(\mathcal{C}^L) = \iota(\tilde{\gamma}_i, \tilde{\beta}_j^L) = 0, \quad (69)$$

$$m_{i,j}(\mathcal{C}^R) = \iota(\tilde{\gamma}_i, \tilde{\beta}_j^R) = 1. \quad (70)$$

It follows that

$$\frac{1}{2}(m_{i,j}(\mathcal{C}^L) + m_{i,j}(\mathcal{C}^R)) = \frac{1}{2} = [\gamma_i^- \gamma_i^+, \eta_j^- \eta_{j+1}^-].$$

We illustrate that case in Figure 7b. This finishes the proof of Equation (60).

Proof of Equation (61) and (62): The proof follows from the same ideas as the previous ones.

□

8. ASYMPTOTICS

This section is the main computational core of this article. Our goal is to compute asymptotic product formulas, namely understand the behavior of the special product formula when the repetition in the arcs becomes infinite. This allows us to describe the limit of certain Wilson loops as elementary functions – see Proposition 8.2.5.

The goal of this section is to obtain Corollary 8.3.2 which is an asymptotic product formula for the Goldman bracket of elementary functions.

We first need some facts about vanishing sequences

8.1. Properties of vanishing sequences. In this paragraph, we shall be given a vanishing sequence $\{\Gamma_p\}_{p \in \mathbb{N}}$ of finite index subgroups of $\pi_1(S)$. We need some notation and definition

- Let gh_p be the function defined on $H(n, S)$ by

$$\text{gh}_p(\rho) = \text{gh}(\rho|_{\Gamma_p}).$$

- For any positive integer p and primitive element ξ in Γ_0 , let $\xi(p)$ be the positive integer so that

$$\langle \xi^{\xi(p)} \rangle = \langle \xi \rangle \cap \Gamma_p.$$

We write $\xi_p = \xi^{\xi(p)}$ and we denote the associated closed geodesic by $\tilde{\xi}_p$.

Definition 8.1.1. [N-NICE COVERING] *Let γ and η be primitive coprime elements of $\Gamma_0 = \pi_1(S)$. Let N be a positive integer. We say that Γ_p is N -nice with respect to γ and η , if the intersection loop $c_x(\tilde{\gamma}_p, \tilde{\eta}_p)$ is either trivial or satisfies*

$$\pi_p(c_x(\tilde{\gamma}_p, \tilde{\eta}_p)) = \gamma^{k_1} \cdot \eta^{-k_2},$$

where k_1 and k_2 satisfy

$$\gamma(p) - N > k_1 > N \text{ and } \eta(p) - N > k_2 > N.$$

We need the following properties of vanishing sequences which we summarize in the next proposition.

Proposition 8.1.2. *Let $\{\Gamma_p\}_{p \in \mathbb{N}}$ be a vanishing sequence of finite index subgroups of $\pi_1(S)$. Then*

- (1) *when p goes to infinity, gh_p converge uniformly to 0 on every compact of $H(n, S)$,*
- (2) *for any primitive coprime elements γ and η , there exists p_0 so that for every $p > p_0$, Γ_p is N -nice with respect to γ and η .*
- (3) *Let $G = (\gamma_0, \dots, \gamma_p)$ and $F = (\eta_0, \dots, \eta_q)$ be tuples of primitive elements of $\pi_1(S) \setminus \{1\}$ such that (γ_i, γ_{i+1}) as well as (η_j, η_{j+1}) are pairwise coprime. Then for p large enough, G and F satisfy the Good Position Hypothesis 7.5.1 as elements of $\pi_1(S_p)$.*

8.1.1. *Proof of Proposition 8.1.2.* Proposition 8.1.2 is the concatenation of Propositions 8.1.3, 8.1.5 and 8.1.6. Proposition 8.1.4 is an intermediate step in proving Proposition 8.1.6. We fix in this paragraph a vanishing sequence $\{\Gamma_p\}_{p \in \mathbb{N}}$.

Remember that we identify primitive elements in $\pi_1(S)$ and in any of its finite index subgroup.

Proposition 8.1.3. *When p goes to infinity, gh_p converge uniformly to 0 on every compact of $\mathbf{H}(n, S)$.*

Proof. For all positive number K and compact C , let

$$S_K := \{\gamma \in \pi_1(S) \setminus \{\text{Id}\} \mid \exists \rho \in C, \text{ width}(\rho(\gamma)) < K\}.$$

Then the set of conjugacy classes in S_K is a finite set. Let S_K^0 be a finite set in $\pi_1(S)$ of representatives of the conjugacy classes of S_K . From the definition of vanishing sequences, it follows that there exists n_0 that that for all $p > p_0$, we have

$$S_K^0 \cap \Gamma_p = \emptyset.$$

Since Γ_p is normal, it follows that

$$S_K \cap \Gamma_p = \emptyset.$$

Then by definition, the girth of any representation in C restricted to Γ_n is greater than K . Thus the family of functions gh_p converges uniformly to zero, when p goes to ∞ . \square

The following proposition is well known.

Proposition 8.1.4. *Let γ be an element of Γ_0 . Then there exists p_0 such that for all $p > p_0$, the geodesics $\tilde{\gamma}_p$ is simple.*

Proof. Let

$$\hat{A}_\gamma := \{\xi \in \Gamma_0 \mid \xi(\tilde{\gamma}) \cap \tilde{\gamma} \neq \emptyset\} \subset \Gamma_0 / \langle \gamma \rangle.$$

Observe that A_γ invariant by right multiplication by γ and that its projection in $\Gamma_0 / \langle \gamma \rangle$ is a finite set. Thus there exists p_0 so that for every $p > p_0$,

$$A_\gamma \cap \Gamma_p \subset \langle \gamma \rangle.$$

This implies that the projection of $\tilde{\gamma}$ in S_p is a simple closed geodesic, indeed the existence of a self intersection point implies the existence of an element ξ in Γ_p so that $\xi(\tilde{\gamma}) \cap \tilde{\gamma} \neq \emptyset$. \square

We finally need.

Proposition 8.1.5. *Let γ and η two coprime primitive elements of $\Gamma_0 = \pi_1(S)$. Let N be a positive integer.*

Then there exists p_0 such that for all $p > p_0$, Γ_p is N -nice with respect to γ and η .

Proof. We assume using the previous proposition that $\tilde{\gamma}_p$ and $\tilde{\eta}_p$ are simple.

STEP 1 We shall prove the following assertion

For any $N > 0$, there exists p_0 such that for any $p > p_0$, for any integers k such that $0 < k \leq N$ and for any m then

$$\gamma^k \eta^m \notin \Gamma_p \text{ and } \gamma^m \eta^{-k} \notin \Gamma_p.$$

This is an immediate application of Proposition 6.2.2. Let $H := \{\gamma^k \mid 0 < k \leq N\}$. Since γ and η are coprime then $H \cap \langle \eta \rangle = \emptyset$. Using Proposition, 6.2.2, we get that they exists p_0 so that for all $p > p_0$,

$$H \cap (\Gamma_p \cdot \langle \eta \rangle) = \emptyset.$$

In other words, for all n and k so that $0 < k \leq N$,

$$\gamma^k \cdot \eta^n \notin \Gamma_p.$$

A symmetric argument concludes the proof.

STEP 2 We now prove.

If $x \in \gamma_p \cap \eta_p$, then there exists positive integers k_1 and k_2 such that the intersection loop $c_p(x) := c_x(\tilde{\gamma}_p, \tilde{\eta}_p)$ satisfies

$$\pi_p(c_p(x)) = \gamma^{k_2} \cdot \eta^{-k_1},$$

where the equality is as homotopy classes in S_0

We can as well assume (using the first step and a shift in p) that the projection of the axis of γ and η are simple geodesics in S_0 . Let also

$$A_p := \{\xi \in \Gamma_p \mid \xi(\tilde{\eta}) \cap \tilde{\gamma} \neq \emptyset\} \subset \Gamma_p.$$

Observe that A_p is invariant by left multiplication by γ and right multiplication by η . Let \hat{A}_p be the projection of A_p in $\langle \eta_p \rangle \backslash \Gamma_p / \langle \eta_p \rangle$. Observe also that we have a bijection from \hat{A}_p to

$$I_p := \pi_p(\tilde{\gamma}) \cap \pi_p(\tilde{\eta}) \subset S_p$$

given by

$$\langle \gamma \rangle \cdot \xi \cdot \langle \eta \rangle \rightarrow \pi_p(\xi(\tilde{\eta}) \cap \tilde{\gamma}).$$

In particular \hat{A}_p is finite since I_p is finite. Moreover, if x in I_p comes in this procedure from an element a in A_p , then a represents the intersection loop of x .

Since \hat{A}_0 is finite, using the double coset separability property, there exists p_0 such that for all $p > p_0$, we have

$$A_0 \cap \Gamma_p \subset \langle \gamma \rangle \langle \eta \rangle,$$

Since $A_p \subset A_0 \cap \Gamma_p$, It follows that the projection in S_0 of any intersection loop $c_x(\gamma_p, \eta_p)$ is homotopic to $\gamma^n \cdot \eta^{-m}$ with n and m positive integers.

CONCLUSION OF THE PROOF The Proposition follows at once from the two steps of the proof. \square

Proposition 8.1.6. *Let $G = (\gamma_0, \dots, \gamma_p)$ and $F = (\eta_0, \dots, \eta_q)$ be tuples of primitive elements of $\pi_1(S) \setminus \{1\}$ such that (γ_i, γ_{i+1}) as well as (η_j, η_{j+1}) are pairwise coprime. Then for p large enough, G and F satisfy the Good Position Hypothesis 7.5.1 as elements of $\pi_1(S_p)$.*

Proof. Let us check the three conditions of the Good Position Hypothesis. Let $G = (\gamma_0, \dots, \gamma_p)$ and $F = (\eta_0, \dots, \eta_q)$ be primitive elements of $\pi_1(S) \setminus \{1\}$ such that (γ_i, γ_{i+1}) as well as (η_j, η_{j+1}) are pairwise coprime.

- (1) Let $B \subset \tilde{S}$ be ball containing all the intersections $\tilde{\gamma}_i \cap \tilde{\eta}_j$ when γ_i and η_j are coprime. Thus Condition (1) of the Good Position Hypothesis is satisfied.
- (2) Let

$$F := \{\xi \in \pi_1(S) \mid B \cap \xi(B) \neq \emptyset\}.$$

The set F is finite. Thus, by Proposition 6.2.2 (applied to $\gamma = \eta = \text{Id}$), there exists p_0 , such that for all $p > p_0$, we have

$$F \cap \Gamma_p = \{\text{Id}\}.$$

Thus Condition (2) of the Good Position Hypothesis is satisfied.

- (3) Next, for every $\zeta \in \{\gamma_0, \dots, \gamma_p, \eta_0, \dots, \eta_q\}$ the set

$$H_\zeta := \{\xi \in \Gamma / \langle \zeta \rangle \mid \xi(\tilde{\zeta}) \cap B \neq \emptyset\}.$$

finite. Thus by Proposition 6.2.2 (applied to $\gamma = \text{Id}$, $\eta = \zeta$), there exists p_0 , such that for all $p > p_0$, we have

$$H_\zeta \langle \zeta \rangle \cap \Gamma_p = \langle \zeta \rangle.$$

Thus Condition (3) of the Good Position Hypothesis is satisfied.

- (4) Finally Condition (4) of the Good Position Hypothesis is satisfied for p large enough by Proposition 8.1.4.

\square

8.2. Asymptotic product formula for Wilson loops. In all this paragraph, we shall be given a finite index subgroup Γ_k of $\Gamma_0 = \pi_1(S)$, corresponding to a covering $S_k \rightarrow S_0 = S$. Then, if ρ is a Hitchin representation of $\pi_1(S)$ in $\mathrm{PSL}_n(\mathbb{R})$, ρ_k will denote the restriction of ρ to Γ_k .

Let $(\gamma_0, \dots, \gamma_q)$ and $(\eta_0, \dots, \eta_{q'})$ be two tuples of primitive elements of $\pi_1(S)$. We assume that (γ_i, γ_{i+1}) as well as (η_j, η_{j+1}) are all pairwise coprime.

Let then $\hat{\gamma}_i$ and $\hat{\eta}_i$ be the representatives of γ_i and η_i in Γ_k , and

$$\mathbf{F}^{(p)} = \hat{\gamma}_1^p \dots \hat{\gamma}_q^p, \quad \mathbf{G}^{(p)} = \hat{\eta}_1^p \dots \hat{\eta}_{q'}^p. \quad (71)$$

We want to understand the asymptotics when p goes to infinity of the following function

$$B_p^k(\gamma_0, \dots, \gamma_q; \eta_0, \dots, \eta_{q'}) : \mathbf{H}(n, S_k) \rightarrow \mathbb{R},$$

defined by

$$B_p^k(\gamma_0, \dots, \gamma_q; \eta_0, \dots, \eta_{q'}) := \frac{\mathbf{W}(\{\mathbf{G}^{(p)}, \mathbf{F}^{(p)}\}_{S_k})}{\mathbf{W}(\mathbf{G}^{(p)})\mathbf{W}(\mathbf{F}^{(p)})} \quad (72)$$

Let then

$$\begin{aligned} \mathbf{f}_{i,j} &= [\gamma_i^- \gamma_i^+, \eta_j^- \eta_j^+], & \mathbf{n}_{i,j} &= [\gamma_i^- \gamma_{i+1}^-, \eta_j^- \eta_j^+], \\ \mathbf{m}_{i,j} &= [\gamma_i^- \gamma_i^+, \eta_j^- \eta_{j+1}^-], & \mathbf{q}_{i,j} &= [\gamma_i^- \gamma_{i+1}^-, \eta_j^- \eta_{j+1}^-]. \end{aligned} \quad (73)$$

The following paragraph is devoted to the proof of the following Proposition

Proposition 8.2.1. [ASYMPTOTIC PRODUCT FORMULA] *For every compact set U in $\mathbf{H}(n, S)$, for every positive integer N , then for k large enough, we have for every ρ in U*

$$\begin{aligned} B_p^k(\gamma_0, \dots, \gamma_q, \eta_0, \dots, \eta_{q'}) (\rho) &= \sum_{\substack{0 \leq i \leq q \\ 0 \leq j \leq q'}} \left(p \mathbf{f}_{i,j} + (p-1)^2 \mathbf{f}_{i,j} \mathbf{T}(\gamma_i, \eta_j) \right. \\ &+ (p-1) \left(\frac{\mathbf{T}(\gamma_{i+1}, \gamma_i, \eta_j)}{\mathbf{T}(\gamma_i, \gamma_{i+1})} (\mathbf{n}_{i,j} + \mathbf{f}_{i+1,j}) + \frac{\mathbf{T}(\gamma_i, \eta_{j+1}, \eta_j)}{\mathbf{T}(\eta_j, \eta_{j+1})} (\mathbf{m}_{i,j} + \mathbf{f}_{i,j+1}) \right) \\ &\left. + \frac{\mathbf{T}(\gamma_{i+1}, \gamma_i, \eta_{j+1}, \eta_j)}{\mathbf{T}(\gamma_{i+1}, \gamma_i) \mathbf{T}(\eta_j, \eta_{j+1})} (\mathbf{q}_{i,j} + \mathbf{n}_{i,j+1} + \mathbf{m}_{i+1,j} + \mathbf{f}_{i+1,j+1}) \right) \\ &\quad p^2 \mathbf{R}_{i,j} \mathbf{T}(\gamma_i, \eta_j) + K(\mathrm{gh}_k(\rho) + \mathrm{gh}_0(\rho)^N), \end{aligned} \quad (74)$$

where

- K is bounded on U ,

- $\text{gh}(\rho)$ is the girth of ρ as defined in Definition 4.1.2
- the integers $f_{i,j}$, $m_{i,j}$, $n_{i,j}$ and $q_{i,j}$ are defined in Equations (73).
- $R_{i,j}$ is an integer that only depends on γ_i and η_j .

We will use bouquets to express this asymptotics using our product formula for bouquets.

8.2.1. *Preliminary asymptotics.* Let ρ be a representation of $\Gamma_0 = \pi_1(S)$. For any k , let $\rho_k := \rho|_{\Gamma_k}$. Let $\gamma_0, \dots, \gamma_q$ and $\eta_0, \dots, \eta_{q'}$ be primitive elements of Γ_0 and $\widehat{\gamma}_0, \dots, \widehat{\gamma}_q$ and $\widehat{\eta}_0, \dots, \widehat{\eta}_{q'}$ be the corresponding elements in Γ_k , so that

$$\widehat{\gamma}_i = \gamma_i^{Q_i}, \quad \widehat{\eta}_j = \eta_j^{P_j}, \quad (75)$$

where Q_i and P_j are positive integers. In this proof, K, K_0, K_1, \dots will be the generic symbol for a function of bounded by a continuous function that only depends on the relative position of the eigenvectors of $\rho(\gamma_i)$ and $\rho(\eta_i)$ and does not depend on k . Let us define

$$\widehat{\mathbf{g}}_i = \rho_k(\widehat{\gamma}_i), \quad \widehat{\mathbf{h}}_i = \rho_k(\widehat{\eta}_i), \quad (76)$$

$$\mathbf{g}_i = \rho(\gamma_i), \quad \mathbf{h}_i = \rho(\eta_i), \quad (77)$$

and

$$\widehat{\mathbf{F}}_i^{(p,m)} := \widehat{\mathbf{g}}_i^n \widehat{\mathbf{g}}_{i+1}^p \dots \widehat{\mathbf{g}}_{i-1}^p \widehat{\mathbf{g}}_i^{p-n}, \quad (78)$$

$$\widehat{\mathbf{G}}_i^{(p,m)} := \widehat{\mathbf{h}}_i^n \widehat{\mathbf{h}}_{i+1}^p \dots \widehat{\mathbf{h}}_{i-1}^p \widehat{\mathbf{h}}_i^{p-n}. \quad (79)$$

$$\mathbf{F}_i^{(p,m)} := \mathbf{g}_i^n \mathbf{g}_{i+1}^p \dots \mathbf{g}_{i-1}^p \mathbf{g}_i^{p-n}, \quad (80)$$

$$\mathbf{G}_i^{(p,m)} := \mathbf{h}_i^n \mathbf{h}_{i+1}^p \dots \mathbf{h}_{i-1}^p \mathbf{h}_i^{p-n}. \quad (81)$$

We prove in this paragraph two propositions

Proposition 8.2.2. *For all positive integer p , for all integer m , with $0 < m < p$, we have for any ρ in a compact set U of $\mathbf{H}(n, S)$,*

$$\frac{\widehat{\mathbf{F}}_i^{(p,0)}}{\text{tr}(\widehat{\mathbf{F}}_i^{(p,0)})} = \frac{\dot{\mathbf{g}}_i \dot{\mathbf{g}}_{i-1}}{\text{tr}(\dot{\mathbf{g}}_i \dot{\mathbf{g}}_{i-1})} + K_1 \cdot \text{gh}_k(\rho)^p, \quad (82)$$

$$\frac{\widehat{\mathbf{F}}_i^{(p,p)}}{\text{tr}(\widehat{\mathbf{F}}_i^{(p,p)})} = \frac{\dot{\mathbf{g}}_{i+1} \dot{\mathbf{g}}_i}{\text{tr}(\dot{\mathbf{g}}_{i+1} \dot{\mathbf{g}}_i)} + K_2 \cdot \text{gh}_k(\rho)^p, \quad (83)$$

$$\frac{\widehat{\mathbf{F}}_i^{(p,m)}}{\text{tr}(\widehat{\mathbf{F}}_i^{(p,m)})} = \dot{\mathbf{g}}_i + K_3 \cdot \text{gh}_k(\rho)^{\inf(m, m-p)}, \quad (84)$$

where K_i are locally bounded functions of ρ .

We recall that $\dot{\mathbf{g}}$ is the projector on the eigendirection of the highest eigenvalue of \mathbf{g} .

Proof. We use Proposition 4.1.4. Thus for all $p \neq m > 0$, we have

$$\widehat{\mathbf{F}}_i^{(p,m)} = \left(\prod_{j=0}^{j=q} \mathbf{W}(\widehat{\gamma}_j)^p \right) (\dot{\mathbf{g}}_i \dot{\mathbf{g}}_{i+1} \dots \dot{\mathbf{g}}_{i-1} \dot{\mathbf{g}}_i + K_0 \cdot \text{gh}_k(\rho)^{\inf(m,m-p)}) . \quad (85)$$

Similarly for $n = 0$ or $n = p$, we have

$$\begin{aligned} \widehat{\mathbf{F}}_i^{(p,0)} &= \left(\prod_{j=0}^{j=q} \mathbf{W}(\widehat{\gamma}_j)^p \right) (\dot{\mathbf{g}}_{i+1} \dots \dot{\mathbf{g}}_{i-1} \dot{\mathbf{g}}_i + K_1 \cdot \text{gh}_k(\rho)^p) , \\ \widehat{\mathbf{F}}_i^{(p,p)} &= \left(\prod_{j=0}^{j=q} \mathbf{W}(\widehat{\gamma}_j)^p \right) (\dot{\mathbf{g}}_i \dot{\mathbf{g}}_{i+1} \dots \dot{\mathbf{g}}_{i-1} + K_2 \cdot \text{gh}_k(\rho)^p) , \end{aligned} \quad (86)$$

Observe now that if $\widehat{\mathbf{p}}$ and $\widehat{\mathbf{q}}$ are projectors such that $\text{tr}(\mathbf{p}, \mathbf{q}) \neq 0$ then for every endomorphism A ,

$$\mathbf{p}A\mathbf{q} = \frac{\text{tr}(\mathbf{p}A\mathbf{q})}{\text{tr}(\mathbf{p}\mathbf{q})} \mathbf{p}\mathbf{q}.$$

Thus for all p

$$\frac{\widehat{\mathbf{F}}_i^{(p,0)}}{\text{tr}(\widehat{\mathbf{F}}_i^{(p,0)})} = \frac{\dot{\mathbf{g}}_i \dot{\mathbf{g}}_{i-1}}{\text{tr}(\dot{\mathbf{g}}_i \dot{\mathbf{g}}_{i-1})} + K_3 \cdot \text{gh}_k(\rho)^p, \quad (87)$$

$$\frac{\widehat{\mathbf{F}}_i^{(p,p)}}{\text{tr}(\widehat{\mathbf{F}}_i^{(p,p)})} = \frac{\dot{\mathbf{g}}_{i+1} \dot{\mathbf{g}}_i}{\text{tr}(\dot{\mathbf{g}}_{i+1} \dot{\mathbf{g}}_i)} + K_4 \cdot \text{gh}_k(\rho)^p, \quad (88)$$

$$\frac{\widehat{\mathbf{F}}_i^{(p,m)}}{\text{tr}(\widehat{\mathbf{F}}_i^{(p,m)})} = \dot{\mathbf{g}}_i + K_5 \cdot \text{gh}_k(\rho)^{\inf(m,m-p)}, \text{ for } m \notin \{0, p\}. \quad (89)$$

□

We use the same notation as in the beginning of the paragraph.

Proposition 8.2.3. *Let us fix i and j . Let*

- $\{N_1, \dots, N_r\}$ *be an sequence of pairwise distinct integers so that $N_l \geq N$ and $Q_j - N_l \geq N$.*
- $\{M_1, \dots, M_r\}$ *be an sequence of pairwise distinct integers so that $M_l \geq N$ and $P_j - M_l \geq N$.*

Then, we have for any ρ in a compact set U in $H(n, S)$, for any positive integers p, q, m and m' ,

$$\sum_{1 \leq l \leq r} \frac{\mathbf{g}_i^{-N_l} \widehat{\mathbf{F}}_i^{(p,m)} \mathbf{g}_i^{N_l} \cdot \mathbf{h}_j^{-M_l} \widehat{\mathbf{G}}_j^{(p,m')} \mathbf{g}_i^{M_l}}{\mathrm{tr} \left(\widehat{\mathbf{F}}_i^{(p,0)} \right) \mathrm{tr} \left(\widehat{\mathbf{G}}_j^{(p,0)} \right)} = r \cdot \mathbf{g}_i \cdot \mathbf{h}_j + K \cdot \mathrm{gh}_0(\rho)^{M+N}, \quad (90)$$

where K is a locally bounded function of ρ and

$$M = \inf(Q_i(m-1), P_j(m'-1), Q_i p - Q m', M_j p - m)$$

Proof. In this proof as usual K_i will denote a locally bounded function of ρ . For the purpose of this proof, we define

$$\widetilde{F}_i^{(p)} = \widehat{\mathbf{g}}_{i+1}^p \cdots \widehat{\mathbf{g}}_{i-1}^p, \quad \widetilde{G}_j^{(p)} = \widehat{\mathbf{h}}_{j+1}^p \cdots \widehat{\mathbf{h}}_{j-1}^p.$$

By definition, if $m \geq 1$, $m' \geq 1$, $n < Q_i$ and $r < P_j$,

$$\begin{aligned} \mathbf{g}_i^{-n} \widehat{\mathbf{F}}_i^{(p,m)} \mathbf{g}_i^n &= \mathbf{g}_i^{Q_i m - n} \widetilde{F}_i^{(p)} \mathbf{g}_i^{Q_i(p-m)+n}, \\ \mathbf{h}_j^{-r} \widehat{\mathbf{G}}_j^{(p,m')} \mathbf{h}_j^r &= \mathbf{h}_j^{P_j m' - r} \widetilde{G}_j^{(p)} \mathbf{h}_j^{P_j(p-m')+r}. \end{aligned}$$

Observe also that

$$\begin{aligned} \mathrm{tr} \left(\widehat{\mathbf{F}}_i^{(p,0)} \right) &= \mathrm{tr} \left(\widehat{\mathbf{F}}_i^{(q,m)} \right), \\ \mathrm{tr} \left(\widehat{\mathbf{G}}_j^{(p,0)} \right) &= \mathrm{tr} \left(\widehat{\mathbf{G}}_j^{(p,m')} \right). \end{aligned}$$

Thus using the asymptotics of Proposition 4.1.4, we get that

$$\frac{\mathbf{g}_i^{-N_l} \widehat{\mathbf{F}}_i^{(p,m)} \mathbf{g}_i^{N_l} \cdot \mathbf{h}_j^{-M_l} \widehat{\mathbf{G}}_j^{(p,m')} \mathbf{g}_i^{M_l}}{\mathrm{tr} \left(\widehat{\mathbf{F}}_i^{(p,0)} \right) \mathrm{tr} \left(\widehat{\mathbf{G}}_j^{(p,0)} \right)} = r \mathbf{g}_i \cdot \mathbf{h}_j + K_0 \cdot \left(\sum_{1 \leq l \leq r} \mathrm{gh}_0(\rho)^{R_l} \right).$$

where

$$R_l := \inf(Q_i m - N_l, Q_i p - Q m + N_l, P_j m' - M_l, P_j p - P m' + M_l).$$

To conclude the proof, we will show that

$$\sum_{1 \leq l \leq r} \mathrm{gh}_0(\rho)^{R_l} \leq \frac{4 \mathrm{gh}_0(\rho)^N}{1 - \mathrm{gh}_0(\rho)}. \quad (91)$$

Let

$$\begin{aligned} \mathcal{A} &= \{l \mid R_l = Q_i m - N_l\}, \\ \mathcal{B} &= \{l \mid R_l = Q_i p - Q m + N_l\}, \\ \mathcal{C} &= \{l \mid R_l = P_j m' - M_l\}, \\ \mathcal{D} &= \{l \mid R_l = P_j p - P m' + M_l\}. \end{aligned}$$

By definition,

$$\begin{aligned} \sum_{l \in \mathcal{A}} \text{gh}_0(\rho)^{R_l} &= \sum_{l \in \mathcal{A}} \text{gh}_0(\rho)^{Q_{m-N_l}} \\ &\leq \sum_{n \geq Q_i(m-1)+N} \text{gh}_0(\rho)^n \\ &\leq \frac{\text{gh}_0(\rho)^{N+Q_i(m-1)}}{1 - \text{gh}_0(\rho)}. \end{aligned}$$

Symmetric arguments show that

$$\begin{aligned} \sum_{i \in \mathcal{B}} \text{gh}_0(\rho)^{R_i} &\leq \frac{\text{gh}_0(\rho)^{N+Q_i(p-m)}}{1 - \text{gh}_0(\rho)} \\ \sum_{l \in \mathcal{C}} \text{gh}_0(\rho)^{R_l} &\leq \frac{\text{gh}_0(\rho)^{N+P_j(m'-1)}}{1 - \text{gh}_0(\rho)} \\ \sum_{l \in \mathcal{D}} \text{gh}_0(\rho)^{R_l} &\leq \frac{\text{gh}_0(\rho)^{N+P_j(p-m')}}{1 - \text{gh}_0(\rho)}. \end{aligned}$$

Inequality (91) – and thus the result – follows. \square

8.2.2. Asymptotics and bouquets. We use the same notation as in the beginning of this section: Let $G = (\gamma_0, \dots, \gamma_q)$ and $(F = \eta_0, \dots, \eta_{q'})$ be two tuples of primitive elements of $\pi_1(S)$. We assume that (γ_i, γ_{i+1}) as well as (η_j, η_{j+1}) are all pairwise coprime. We shall use the notation of Paragraph 7.3.2.

Proposition 8.2.4. *Assume that G and F and Γ_k satisfy the Good Position Hypothesis. Assume also that Γ_k is N -nice for all pairs (γ_i, η_j) . Let C be a bouquet in a good position representing G and F .*

Then for every compact set U in $\mathbf{H}(n, S)$, we have for every ρ in U ,

$$\begin{aligned} B_p^k(\gamma_0, \dots, \gamma_q, \eta_0, \dots, \eta_{q'})(&\rho) = \sum_{\substack{0 \leq i \leq q \\ 0 \leq j \leq q'}} \left((p-1)^2 \mathbf{f}_{i,j}(\mathcal{C}) \mathbf{T}(\gamma_i, \eta_j) \right. \\ &+ (p-1) \left(\frac{\mathbf{T}(\gamma_{i+1}, \gamma_i, \eta_j)}{\mathbf{T}(\gamma_i, \gamma_{i+1})} (\mathbf{n}_{i,j}(\mathcal{C}) + \mathbf{f}_{i+1,j}(\mathcal{C})) + \frac{\mathbf{T}(\gamma_i, \eta_{j+1}, \eta_j)}{\mathbf{T}(\eta_j, \eta_{j+1})} (\mathbf{m}_{i,j}(\mathcal{C}) + \mathbf{f}_{i,j+1}(\mathcal{C})) \right) \\ &+ \frac{\mathbf{T}(\gamma_{i+1}, \gamma_i, \eta_{j+1}, \eta_j)}{\mathbf{T}(\gamma_{i+1}, \gamma_i) \mathbf{T}(\eta_j, \eta_{j+1})} (\mathbf{q}_{i,j}(\mathcal{C}) + \mathbf{n}_{i,j+1}(\mathcal{C}) + \mathbf{m}_{i+1,j}(\mathcal{C}) + \mathbf{f}_{i+1,j+1}(\mathcal{C})) \Big) \\ &+ p^2 \left(\sum_{i,j} I_{i,j}(1) \sharp(C_{i,j}) \mathbf{T}(\gamma_i, \eta_j) \right) + K(\text{gh}_k(\rho) + \text{gh}_0(\rho)^N), \quad (92) \end{aligned}$$

where

- K is bounded by a continuous function that only depends on the relative position of the eigenvectors of $\rho(\gamma_i)$ and $\rho(\eta_j)$.
- $\text{gh}(\rho)$ is the girth of ρ as defined in Definition 4.1.2
- the integers $f_{i,j}(\mathcal{C})$, $m_{i,j}(\mathcal{C})$, $n_{i,j}(\mathcal{C})$ and $q_{i,j}(\mathcal{C})$ are defined in Equations (53).

8.2.3. *Proof of Proposition 8.2.5.* We now recall the the Product Formula (54) that we write using the Notation of Paragraph 8.2.1 as

$$B_p = B_p^0 + \sum_{\xi \in C_{i,j}} I_{i,j}(\xi) B_p^\xi, \quad (93)$$

where

$$\begin{aligned} B_p^0 := & \frac{1}{\text{tr}(\widehat{F}^{(p,0)}) \cdot \text{tr}(\widehat{G}^{(p,0)})} \sum_{\substack{0 \leq i \leq q \\ 0 \leq j \leq q'}} \left(\sum_{\substack{1 \leq m' \leq p \\ 1 \leq m \leq p}} f_{i,j}(\mathcal{C}) \text{tr}(\widehat{F}_i^{(p,m')} \widehat{G}_j^{(p,m)}) \right. \\ & + \sum_{1 \leq m \leq p} m_{i,j}(\mathcal{C}) \text{tr}(\widehat{F}_i^{(p,m)} \widehat{G}_j^{(p,0)}) \\ & \left. + \sum_{1 \leq m' \leq p} n_{i,j}(\mathcal{C}) \text{tr}(\widehat{F}_i^{(p,0)} \widehat{G}_j^{(p,m')}) + q_{i,j}(\mathcal{C}) \text{tr}(\widehat{F}_i^{(p,0)} \widehat{G}_j^{(p,0)}) \right) \quad (94) \end{aligned}$$

and

$$B_p^\xi := \frac{1}{\text{tr}(\widehat{F}^{(p,0)}) \cdot \text{tr}(\widehat{G}^{(p,0)})} \sum_{\substack{1 \leq m' \leq p \\ 1 \leq m \leq p}} \text{tr}(\widehat{F}_i^{(p,m')} \rho(\xi) \widehat{G}_j^{(p,m)} \rho(\xi)^{-1}) \quad (95)$$

Proposition 8.2.5 will follow from the two next propositions that treat independently the term B_p^0 and the term involving the B_p^ξ .

Proposition 8.2.5. *We have*

$$\begin{aligned} B_p^0(\gamma_0, \dots, \gamma_q, \eta_0, \dots, \eta_{q'}) = & \sum_{\substack{0 \leq i \leq q \\ 0 \leq j \leq q'}} \left((p-1)^2 f_{i,j}(\mathcal{C}) \text{T}(\gamma_i, \eta_j) \right. \\ & + (p-1) \left(\frac{\text{T}(\gamma_{i+1}, \gamma_i, \eta_j)}{\text{T}(\gamma_i, \gamma_{i+1})} (n_{i,j}(\mathcal{C}) + f_{i+1,j}(\mathcal{C})) + \frac{\text{T}(\gamma_i, \eta_{j+1}, \eta_j)}{\text{T}(\eta_j, \eta_{j+1})} (m_{i,j}(\mathcal{C}) + f_{i,j+1}(\mathcal{C})) \right) \\ & \left. + \frac{\text{T}(\gamma_{i+1}, \gamma_i, \eta_{j+1}, \eta_j)}{\text{T}(\gamma_{i+1}, \gamma_i) \text{T}(\eta_j, \eta_{j+1})} (q_{i,j}(\mathcal{C}) + n_{i,j+1}(\mathcal{C}) + m_{i+1,j}(\mathcal{C}) + f_{i+1,j+1}(\mathcal{C})) \right) + K \cdot \text{gh}_k(\rho), \quad (96) \end{aligned}$$

where K only depends on the position of the eigenvectors of $\rho(\gamma_i)$ and $\rho(\eta_j)$.

Proof. Using the estimates for $\widehat{F}_i^{(p,m)}$ and $\widehat{G}_i^{(p,m)}$ coming from Proposition 8.2.2 we get that

$$\begin{aligned}
 B_p^0 = & \sum_{\substack{0 \leq i \leq q \\ 0 \leq j \leq q'}} \left(f_{i,j}(\mathcal{C})(p-1)^2 \operatorname{tr}(\dot{\mathbf{g}}_i \dot{\mathbf{h}}_j) \right. \\
 & + f_{i,j}(\mathcal{C}) \left(\frac{\operatorname{tr}(\dot{\mathbf{g}}_i \dot{\mathbf{g}}_{i-1} \dot{\mathbf{h}}_j \dot{\mathbf{h}}_{j-1})}{\operatorname{tr}(\dot{\mathbf{g}}_i \dot{\mathbf{g}}_{i-1}) \operatorname{tr}(\dot{\mathbf{h}}_j \dot{\mathbf{h}}_{j-1})} + (p-1) \left(\frac{\operatorname{tr}(\dot{\mathbf{g}}_i \dot{\mathbf{h}}_j \dot{\mathbf{h}}_{j-1})}{\operatorname{tr}(\dot{\mathbf{h}}_j \dot{\mathbf{h}}_{j-1})} + \frac{\operatorname{tr}(\dot{\mathbf{h}}_j \dot{\mathbf{g}}_i \dot{\mathbf{g}}_{i-1})}{\operatorname{tr}(\dot{\mathbf{g}}_i \dot{\mathbf{g}}_{i-1})} \right) \right) \\
 & + m_{i,j}(\mathcal{C}) \left((p-1) \frac{\operatorname{tr}(\dot{\mathbf{g}}_i \dot{\mathbf{h}}_{j+1} \dot{\mathbf{h}}_j)}{\operatorname{tr}(\dot{\mathbf{h}}_j \dot{\mathbf{h}}_{j+1})} + \frac{\operatorname{tr}(\dot{\mathbf{g}}_i \dot{\mathbf{g}}_{i-1} \dot{\mathbf{h}}_{j+1} \dot{\mathbf{h}}_j)}{\operatorname{tr}(\dot{\mathbf{g}}_i \dot{\mathbf{g}}_{i-1}) \operatorname{tr}(\dot{\mathbf{h}}_{j+1} \dot{\mathbf{h}}_j)} \right) \\
 & + n_{i,j}(\mathcal{C}) \left((p-1) \frac{\operatorname{tr}(\dot{\mathbf{g}}_{i+1} \dot{\mathbf{g}}_i \dot{\mathbf{h}}_j)}{\operatorname{tr}(\dot{\mathbf{g}}_{i+1} \dot{\mathbf{g}}_i)} + \frac{\operatorname{tr}(\dot{\mathbf{g}}_{i+1} \dot{\mathbf{g}}_i \dot{\mathbf{h}}_j \dot{\mathbf{h}}_{j-1})}{\operatorname{tr}(\dot{\mathbf{g}}_i \dot{\mathbf{g}}_{i-1}) \operatorname{tr}(\dot{\mathbf{h}}_{j+1} \dot{\mathbf{h}}_j)} \right) \\
 & + q_{i,j}(\mathcal{C}) \frac{\operatorname{tr}(\dot{\mathbf{g}}_{i+1} \dot{\mathbf{g}}_i \dot{\mathbf{h}}_{j+1} \dot{\mathbf{h}}_j)}{\operatorname{tr}(\dot{\mathbf{g}}_{i+1} \dot{\mathbf{g}}_i) \operatorname{tr}(\dot{\mathbf{h}}_{j+1} \dot{\mathbf{h}}_j)} \Big) + K \cdot \operatorname{gh}_k(\rho). \tag{97}
 \end{aligned}$$

Using the definition of multi fractions, and after reordering terms, we obtain the asymptotics of the proposition. \square

Finally we need to understand the last term involving the sum of the terms B_p^ξ .

Proposition 8.2.6. *We have*

$$\sum_{i,j} \sum_{\xi \in C_{i,j}} I_{i,j}(\xi) B_p^\xi = p^2 \left(\sum_{i,j} I_{i,j}(1) \sharp(C_{i,j}) \operatorname{tr}(\dot{\mathbf{g}}_i \dot{\mathbf{h}}_j) \right) + K \cdot \operatorname{gh}(\rho_0)^N, \tag{98}$$

where K only depends on the position of the eigenvectors of $\rho(\gamma_i)$ and $\rho(\eta_j)$.

Proof. We use again the notation set in the beginning of Paragraph 8.2.1. By definition of an N -nice covering, any element $\xi \in C_{i,j}$ can be written as

$$\xi = \gamma_i^{N_\xi} \eta_j^{M_\xi},$$

where $N < N_\xi < Q_i - N$, and $N < M_\xi < P_j - N$. Since for $0 < m < Q_j$, $\gamma_i^m \notin \Gamma_k$, we obtain that $\xi \rightarrow N_\xi$ and $\xi \rightarrow M_\xi$ are bijections.

Moreover, since the bouquet C is a lift of a bouquet C^0 in S_0 , then

$$I_{i,j}(\xi) = I_{i,j}(1).$$

It follows that for any i and j ,

$$\sum_{\xi \in C_{i,j}} I_{i,j}(\xi) B_p^\xi = I_{i,j}(1) \sum_{\substack{1 \leq m' \leq p \\ 1 \leq m \leq p}} \frac{\text{tr}(B_p^{m,m',i,j})}{\text{tr}(\widehat{F}^{(p,0)}) \cdot \text{tr}(\widehat{G}^{(p,0)})} \quad (99)$$

where

$$B_p^{m,m',i,j} = \sum_{\xi \in C_{i,j}} \left(\mathbf{g}_i^{-N_\xi} \widehat{F}_i^{(p,m)} \mathbf{g}_i^{N_\xi} \cdot \mathbf{h}_j^{-M_\xi} \widehat{G}_j^{(p,m')} \mathbf{h}_j^{M_\xi} \right).$$

We now apply Proposition 8.2.3 to get

$$\frac{B_p^{m,m',i,j}}{\text{tr}(\widehat{F}^{(p,0)}) \cdot \text{tr}(\widehat{G}^{(p,0)})} = I_{i,j}(1) \sharp(C_{i,j}) \dot{\mathbf{g}}_i \dot{\mathbf{h}}_j + K_0 \text{gh}_0(\rho)^{N+M(m,m')}, \quad (100)$$

where $M(m, m') = \inf(Q_i(p - m), Q_i(m - 1), P_j(p - m'), P_j(m' - 1))$. Observe that for any $\lambda < 1$

$$\sum_{\substack{1 \leq m' \leq p \\ 1 \leq m \leq p}} \lambda^{M(m,m')} \leq 4 \sum_{n \leq 0} \lambda^n = \frac{4}{1 - \lambda}.$$

Thus Equations (99) and (100) together yield

$$\sum_{i,j} \sum_{\xi \in C_{i,j}} I_{i,j}(\xi) B_p^\xi = p^2 \left(\sum_{i,j} I_{i,j}(1) \sharp(C_{i,j}) \cdot \text{tr}(\dot{\mathbf{g}}_i \dot{\mathbf{h}}_j) \right) + K \text{gh}_0(\rho)^N \quad (101)$$

The result finally follows from the fact that $\text{tr}(\dot{\mathbf{g}}_i \dot{\mathbf{h}}_j) = \mathsf{T}(\gamma_i, \eta_j)$. \square

8.2.4. *Proof of Proposition 8.2.1.* Since G , F and Γ_k satisfy the Good Condition Hypothesis, by Proposition 7.5.2, there exist two bouquets \mathcal{C}_L and \mathcal{C}_R in S in a homotopically good position, both representing G and F such that furthermore

$$\begin{aligned} \frac{1}{2}(\mathbf{f}_{i,j}(\mathcal{C}_L) + \mathbf{f}_{i,j}(\mathcal{C}_R)) &= \mathbf{f}_{i,j} \\ \frac{1}{2}(\mathbf{n}_{i,j}(\mathcal{C}_L) + \mathbf{n}_{i,j}(\mathcal{C}_R)) &= \mathbf{n}_{i,j}, \\ \frac{1}{2}(\mathbf{m}_{i,j}(\mathcal{C}_L) + \mathbf{m}_{i,j}(\mathcal{C}_R)) &= \mathbf{m}_{i,j}, \\ \frac{1}{2}(\mathbf{q}_{i,j}(\mathcal{C}_L) + \mathbf{q}_{i,j}(\mathcal{C}_R)) &= \mathbf{q}_{i,j}. \end{aligned}$$

Thus applying Proposition 8.2.5 twice, once for \mathcal{C}_L and once for \mathcal{C}_R , and taking the half sum, we obtain the final result.

8.3. Asymptotics of brackets of multifractions. The setting of this paragraph is the same as the previous one: we shall be given a finite index subgroup Γ_k of $\Gamma_0 = \pi_1(S)$, corresponding to a covering $S_k \rightarrow S_0 = S$. Then, if ρ will be a Hitchin representation of $\pi_1(S)$ in $\mathrm{PSL}_n(\mathbb{R})$, ρ_k will denote the restriction of ρ to Γ_k .

Let $G = (\gamma_0, \dots, \gamma_q)$ and $F = (\eta_0, \dots, \eta_{q'})$ be two tuples of primitive elements of $\pi_1(S)$. We assume that (γ_i, γ_{i+1}) as well as (η_j, η_{j+1}) are all pairwise coprime. Observe that there exists $M \in \mathbb{N}$ so that for all i and j , $\hat{\gamma} := \gamma_i^M$ and $\hat{\eta} := \eta_j^M$ belong to Γ_k .

Then let

$$\overline{W}_p(\gamma_1, \dots, \gamma_q) := \frac{W(\hat{\gamma}_1^p \dots \hat{\gamma}_q^p)}{\prod_{i=1}^q W(\hat{\gamma}_i^p)},$$

so that

$$T = \lim_{p \rightarrow \infty} \overline{W}_p. \quad (102)$$

Let now

$$A_p := \frac{\{\overline{W}_p(\gamma_0, \dots, \gamma_q), \overline{W}_p(\eta_0, \dots, \eta_{q'})\}_S}{\overline{W}_p(\gamma_0, \dots, \gamma_q) \cdot \overline{W}_p(\eta_0, \dots, \eta_{q'})} \quad (103)$$

Let $F = (\gamma_1, \dots, \gamma_q)$ and $G = (\eta_1, \dots, \eta_{q'})$. We first have

Proposition 8.3.1. *We have*

$$A_p = B_p(F, G) - \sum_i B_p(\gamma_i, G) - \sum_j B_p(F, \eta_j) + \sum_{\substack{0 \leq i \leq q \\ 0 \leq j \leq q'}} B_p(\gamma_i, \eta_j). \quad (104)$$

From this proposition and Proposition 8.2.5, we will deduce the following important corollary

Corollary 8.3.2. *Assume that G and F and Γ_0 satisfy the Good Position Hypothesis. Let k be a positive integer so that Γ_k is N -nice for all pairs (γ_i, η_j) . Then*

$$\begin{aligned} & \frac{\{T(\gamma_0, \dots, \gamma_q), T(\eta_0, \dots, \eta_{q'})\}_{S_k}}{T(\gamma_0, \dots, \gamma_q) \cdot T(\eta_0, \dots, \eta_{q'})} \\ &= \sum_{\substack{0 \leq i \leq q \\ 0 \leq j \leq q'}} \left((q_{i,j} + n_{i,j+1} + m_{i+1,j} + f_{i+1,j+1}) \frac{T(\gamma_{i+1}\gamma_i, \eta_{j+1}, \eta_j)}{T(\gamma_{i+1}, \gamma_i) T(\eta_j, \eta_{j+1})} \right. \\ & \quad - (n_{i,j} + f_{i+1,j}) \frac{T(\gamma_{i+1}\gamma_i, \eta_j)}{T(\gamma_{i+1}, \gamma_i)} - (m_{i,j} + f_{i,j+1}) \frac{T(\gamma_i, \eta_{j+1}, \eta_j)}{T(\eta_j, \eta_{j+1})} \\ & \quad \left. + f_{i,j} \cdot T(\gamma_i, \eta_j) \right) + K \cdot (\mathrm{gh}_k(\rho) + \mathrm{gh}_0(\rho)^N), \end{aligned} \quad (105)$$

where K is bounded by a continuous function that only depends on the relative position of the eigenvectors of $\rho(\gamma_i)$ and $\rho(\eta_i)$

We first prove the corollary, then the proposition in the new paragraph

Proof. We study one by one the terms in the righthand side of the formula of Proposition 8.3.1 using the asymptotics given by Proposition 8.2.5. Let $\varepsilon = \text{gh}_k(\rho) + \text{gh}_0(\rho)^N$. First,

$$\begin{aligned} B_p(\gamma_0, \dots, \gamma_q, \eta_0, \dots, \eta_{q'}) &= \sum_{\substack{0 \leq i \leq q \\ 0 \leq j \leq q'}} \left((p^2 R_{i,j} + (p-1)^2 f_{i,j}) T(\gamma_i, \eta_j) \right. \\ &+ (p-1) \left(\frac{T(\gamma_{i+1}, \gamma_i, \eta_j)}{T(\gamma_i, \gamma_{i+1})} (n_{i,j} + f_{i+1,j}) + \frac{T(\gamma_i, \eta_{j+1}, \eta_j)}{T(\eta_j, \eta_{j+1})} (m_{i,j} + f_{i,j+1}) \right) \\ &\left. + \frac{T(\gamma_{i+1}, \gamma_i, \eta_{j+1}, \eta_j)}{T(\gamma_{i+1}, \gamma_i) T(\eta_j, \eta_{j+1})} (q_{i,j} + n_{i,j+1} + m_{i+1,j} + f_{i+1,j+1}) \right) + K\varepsilon. \end{aligned} \quad (106)$$

Let us now consider the term $B_p(\gamma_i, \eta_0, \dots, \eta_{q'})$, we can apply Formula (106) using the fact that in this case $q_{i,j} = n_{i,j} = 0$.

Thus

$$\begin{aligned} B_p(\gamma_i, \eta_0, \dots, \eta_{q'}) &= \sum_{0 \leq j \leq q'} \left((p^2 R_{i,j} + (p-1)^2 f_{i,j}) T(\gamma_i, \eta_j) \right. \\ &+ (p-1) \left(T(\gamma_i, \eta_j) (f_{i,j}) + \frac{T(\gamma_i, \eta_{j+1}, \eta_j)}{T(\eta_j, \eta_{j+1})} (m_{i,j} + f_{i,j+1}) \right) \\ &\left. + \frac{T(\gamma_i, \eta_{j+1}, \eta_j)}{T(\eta_j, \eta_{j+1})} (m_{i,j} + f_{i,j+1}) \right) + K\varepsilon. \end{aligned} \quad (107)$$

Similarly,

$$\begin{aligned} B_p(\gamma_0, \dots, \gamma_q, \eta_j) &= \sum_{0 \leq i \leq q} \left((p^2 R_{i,j} + (p-1)^2 f_{i,j}) T(\gamma_i, \eta_j) \right. \\ &+ (p-1) \left(\frac{T(\gamma_{i+1}, \gamma_i, \eta_j)}{T(\gamma_i, \gamma_{i+1})} (n_{i,j} + f_{i+1,j}) + T(\gamma_i, \eta_j) (f_{i,j}) \right) \\ &\left. + \frac{T(\gamma_{i+1}, \gamma_i, \eta_j)}{T(\gamma_{i+1}, \gamma_i)} (n_{i,j} + f_{i+1,j}) \right) + K\varepsilon. \end{aligned} \quad (108)$$

Finally,

$$B_p(\gamma_i, \eta_j) = T(\gamma_i, \eta_j) (p^2 R_{i,j} + (p-1)^2 f_{i,j} + 2(p-1) + 1) + K\varepsilon. \quad (109)$$

Thus, using Proposition 8.3.1, in A_p

- the coefficient of $\mathbb{T}(\gamma_i, \eta_j)$ is $f_{i,j}$,
- the coefficient of $\frac{\mathbb{T}(\gamma_{i+1}, \gamma_i, \eta_j)}{\mathbb{T}(\gamma_i, \gamma_{i+1})}$ is $-(n_{i,j} + f_{i+1,j})$,
- the coefficient of $\frac{\mathbb{T}(\gamma_i, \eta_{j+1}, \eta_j)}{\mathbb{T}(\eta_j, \eta_{j+1})}$ is $-(m_{i,j} + f_{i,j+1})$,
- the coefficient of $\frac{\mathbb{T}(\gamma_{i+1}, \gamma_i, \eta_{j+1}, \eta_j)}{\mathbb{T}(\gamma_{i+1}, \gamma_i) \mathbb{T}(\eta_j, \eta_{j+1})}$ is $q_{i,j} + n_{i,j+1} + m_{i+1,j} + f_{i+1,j+1}$.

Finally, we conclude the proof of the corollary by using Formula (102). \square

8.3.1. *Proof of Proposition 8.3.1.* First we use the “logarithmic derivative formula” for the Poisson bracket

$$\frac{\{f \cdot g, h\}_S}{fgh} = \frac{\{f, h\}_S}{fh} + \frac{\{g, h\}_S}{gh}.$$

We obtain

$$\begin{aligned} A_p(F, G) &= \frac{\{W(\gamma_0^p \dots \gamma_q^p), W(\eta_0^p \dots \eta_{q'}^p)\}_S}{W(\gamma_0^p \dots \gamma_q^p) W(\eta_0^p \dots \eta_{q'}^p)} - \sum_{0 \leq i \leq q} \frac{\{W(\gamma_i^p), W(\eta_0^p \dots \eta_{q'}^p)\}_S}{W(\gamma_i^p) W(\eta_0^p \dots \eta_{q'}^p)} \\ &\quad - \sum_{0 \leq j \leq q'} \frac{\{W(\gamma_0^p \dots \gamma_q^p), W(\eta_j^p)\}_S}{W(\gamma_0^p \dots \gamma_q^p) W(\eta_j^p)} + \sum_{\substack{0 \leq i \leq q \\ 0 \leq j \leq q'}} \frac{\{W(\gamma_i^p), W(\eta_j^p)\}_S}{W(\gamma_i^p) W(\eta_j^p)}. \end{aligned} \quad (110)$$

Then, using the definition of Equation (44) expressing the Goldman Poisson bracket of Wilson loops in terms of the bracket of loops in the Goldman Algebra, we get

$$\frac{\{W(\gamma_0^p \dots \gamma_q^p), W(\eta_0^p \dots \eta_{q'}^p)\}_S}{W(\gamma_0^p \dots \gamma_q^p) W(\eta_0^p \dots \eta_{q'}^p)} = B_p(F, G) - \frac{1}{n} \iota(\gamma_0^p \dots \gamma_q^p, \eta_0^p \dots \eta_{q'}^p).$$

The proposition now follows from the fact that

$$\iota(a \cdot b, c) = \iota(a, c) + \iota(b, c),$$

and thus

$$\begin{aligned} \iota(\gamma_0^p \dots \gamma_q^p, \eta_0^p \dots \eta_{q'}^p) &= \\ &= \sum_{0 \leq i \leq q} \iota(\gamma_i^p, \eta_0^p \dots \eta_{q'}^p) + \sum_{0 \leq j \leq q'} \iota(\gamma_0^p \dots \gamma_q^p, \eta_j^p) - \sum_{\substack{0 \leq i \leq q \\ 0 \leq j \leq q'}} \iota(\gamma_i^p, \eta_j^p). \end{aligned} \quad (111)$$

9. GOLDMAN AND SWAPPING ALGEBRAS: PROOFS OF THE MAIN RESULTS

We finally prove the results stated in Section 6. In the course of the proof, we prove the generalized Wolpert formula in Theorem 9.2.1.

9.1. Poisson brackets of elementary functions and the proof of Theorem 6.2.4. By Corollary 4.2.3, the algebra $\mathcal{B}(\mathbf{P})$ of multi fractions is generated by elementary functions. Thus it is enough to prove the theorem when b_0 and b_1 are elementary functions.

Let $G = (\gamma_0, \dots, \gamma_p)$ and $F = (\eta_0, \dots, \eta_{q'})$ be primitive elements of $\pi_1(S)$. We assume that for all i and j , γ_i and γ_{i+1} are coprime, as well as η_i and η_{i+1} .

Let $b_0 = \mathbf{T}(\gamma_0, \dots, \gamma_q)$ and $b_1 = \mathbf{T}(\eta_0, \dots, \eta_{q'})$.

By Proposition 8.1.6, we can assume that G and F satisfy the Good Position Hypothesis for S_k when $k > k_0$ for some n_0 . Let N be a positive integer, we can further assume that $S_k \mapsto S_0$ is N -nice for all pairs (γ_i, η_j) by Proposition 8.1.5 for $k \geq k_0$ and k_0 large enough.

Recall also, using the notation of Proposition 4.3.1, that

$$\begin{aligned} f_{i,j} &= [\gamma_i^- \gamma_i^+, \eta_j^- \eta_j^+] = a_{i,j} \\ q_{i,j} + n_{i,j+1} + m_{i+1,j} + f_{i+1,j+1} &= [\gamma_i^- \gamma_{i+1}^+, \eta_j^- \eta_{j+1}^+] = b_{i,j}, \\ f_{i,j+1} + m_{i,j} &= [\gamma_i^- \gamma_i^+, \eta_j^- \eta_{j+1}^+] = c_{i,j} \\ f_{i+1,j} + n_{i,j} &= [\gamma_i^- \gamma_{i+1}^+, \eta_j^- \eta_j^+] = d_{i,j}. \end{aligned} \quad (112)$$

Thus Corollary 8.3.2 and the computation of the swapping bracket in Proposition 4.3.1 yield

$$\begin{aligned} & \frac{\{\mathbf{T}(\gamma_0, \dots, \gamma_q), \mathbf{T}(\eta_0, \dots, \eta_{q'})\}_{S_k}}{\mathbf{T}(\gamma_0, \dots, \gamma_q) \cdot \mathbf{T}(\eta_0, \dots, \eta_{q'})} \\ &= \frac{\{\mathbf{T}(\gamma_0, \dots, \gamma_q), \mathbf{T}(\eta_0, \dots, \eta_{q'})\}_W}{\mathbf{T}(\gamma_0, \dots, \gamma_q) \cdot \mathbf{T}(\eta_0, \dots, \eta_{q'})} + K \cdot (\text{gh}_k(\rho) + \text{gh}(\rho)^N), \end{aligned} \quad (113)$$

where K is a bounded function that only depends on the eigenvectors of $\rho(\gamma_i)$ and $\rho(\eta_j)$. In particular, there exists a real number K_0 and a compact neighborhood C of ρ_0 so that the previous equality holds with $K \leq K_0$ and ρ in C .

Let ε be a positive real number. By the last assertion in Proposition 8.1.3, we may furthermore choose k_0 so that if $k > k_0$,

$$\text{gh}_k(\rho) \leq \frac{\varepsilon}{2K_0}.$$

Since $\sup\{\text{gh}(\rho) \mid \rho \in C\} < 1$, we may further choose N – and thus k_0 – so that for all ρ in C ,

$$\text{gh}(\rho)^N \leq \frac{\varepsilon}{2K_0}$$

It follows that for all ρ in C , for all $k \geq k_0$, we have

$$\left| \frac{\{\mathsf{T}(\gamma_0, \dots, \gamma_q), \mathsf{T}(\eta_0, \dots, \eta_{q'})\}_{S_k}}{\mathsf{T}(\gamma_0, \dots, \gamma_q) \cdot \mathsf{T}(\eta_0, \dots, \eta_{q'})} - \frac{\{\mathsf{T}(\gamma_0, \dots, \gamma_q), \mathsf{T}(\eta_0, \dots, \eta_{q'})\}_W}{\mathsf{T}(\gamma_0, \dots, \gamma_q) \cdot \mathsf{T}(\eta_0, \dots, \eta_{q'})} \right| \leq \varepsilon. \quad (114)$$

This concludes the proof of Theorem 6.2.4.

9.2. Poisson brackets of length functions. We shall first prove a result of independent interest, namely the value of the Goldman bracket of two length functions of geodesics having exactly one intersection point.

Given a Hitchin representation ρ in $\mathrm{PSL}_n(\mathbb{R})$, or alternatively a rank- n cross ratio \mathbf{b}_ρ , the period – or length – of a conjugacy class γ in $\pi_1(S)$, is given by

$$\ell_\gamma = \log \left(\frac{\lambda_{\max}(\rho(\gamma))}{\lambda_{\min}(\rho(\gamma))} \right) = \log (\mathbf{b}_\rho(\gamma^-, \gamma^+, y, \gamma(y))), \quad (115)$$

for any $y \in \partial_\infty \pi_1(S)$ different from γ^+ and γ^- and where $\lambda_{\max}(A)$ and $\lambda_{\min}(A)$ denotes respectively the eigenvalue with the greatest and smallest modulus of the endomorphism A .

9.2.1. A generalised Wolpert Formula. We have the following extension of Wolpert Formula:

Theorem 9.2.1. [GENERALISED WOLPERT FORMULA] *Let γ and η two closed geodesic with a unique intersection point then the Goldman bracket of the two length functions ℓ_γ and ℓ_η seen as functions on the Hitchin component is*

$$\{\ell_\gamma, \ell_\eta\}_S = \iota(\gamma, \eta) \sum_{\varepsilon, \varepsilon' \in \{-1, 1\}} \varepsilon \cdot \varepsilon' \cdot \mathsf{T}(\gamma^\varepsilon \cdot \eta^{\varepsilon'}), \quad (116)$$

where we recall that

$$\mathsf{T}(\xi, \zeta)(\rho) = \mathbf{b}_\rho(\xi^+, \zeta^+, \zeta^-, \xi^-).$$

Proof. Let us first remark that

$$\ell_\gamma = \lim_{n \rightarrow +\infty} \frac{1}{n} \log (\mathrm{tr}(\rho(\gamma^n)) \mathrm{tr}(\rho(\gamma^{-n}))). \quad (117)$$

Thus assuming that γ and η have a unique intersection point x , whose intersection number is $\iota(\gamma, \eta)$, the Product Formula (54) gives us if $\varepsilon_i \in \{-1, 1\}$,

$$\{\gamma^{\varepsilon \cdot p}, \eta^{\varepsilon' \cdot p}\} = \varepsilon \cdot \varepsilon' \cdot p^2 \cdot \iota(\gamma, \eta) \gamma^{\varepsilon \cdot p} \eta^{\varepsilon' \cdot p}. \quad (118)$$

It follows that

$$\begin{aligned}
& \left\{ \log \left(W(\gamma^p) W(\gamma^{-p}) \right), \log \left(W(\eta^p) W(\eta^{-p}) \right) \right\}_S \\
&= \sum_{\varepsilon, \varepsilon' \in \{-1, 1\}} \varepsilon \cdot \varepsilon' \cdot \iota(\gamma, \eta) \frac{\{W(\gamma^{\varepsilon \cdot p}), W(\eta^{\varepsilon' \cdot p})\}_S}{W(\gamma^{\varepsilon \cdot p}) \cdot W(\eta^{\varepsilon' \cdot p})} \\
&= \sum_{\varepsilon, \varepsilon' \in \{-1, 1\}} p^2 \cdot \varepsilon \cdot \varepsilon' \cdot \iota(\gamma, \eta) \frac{W(\gamma^{\varepsilon \cdot p} \cdot \eta^{\varepsilon' \cdot p})}{W(\gamma^{\varepsilon \cdot p}) \cdot W(\eta^{\varepsilon' \cdot p})} + \frac{1}{n} \iota(\gamma, \eta) \sum_{\varepsilon, \varepsilon' \in \{-1, 1\}} \varepsilon \cdot \varepsilon' \cdot \iota(\gamma, \eta) \quad (19)
\end{aligned}$$

Thus

$$\begin{aligned}
& \lim_{p \rightarrow \infty} \left(\left\{ \frac{1}{p} \log \left(W(\gamma^p) W(\gamma^{-p}) \right), \frac{1}{p} \log \left(W(\eta^p) W(\eta^{-p}) \right) \right\}_S \right) \\
&= \sum_{\varepsilon, \varepsilon' \in \{-1, 1\}} \varepsilon \cdot \varepsilon' \cdot \iota(\gamma, \eta) \lim_{p \rightarrow \infty} \left(\frac{W(\gamma^{\varepsilon \cdot p} \cdot \eta^{\varepsilon' \cdot p})}{W(\gamma^{\varepsilon \cdot p}) \cdot W(\eta^{\varepsilon' \cdot p})} \right) \\
&= \sum_{\varepsilon, \varepsilon' \in \{-1, 1\}} \varepsilon \cdot \varepsilon' \cdot \iota(\gamma, \eta) T(\gamma^\varepsilon \cdot \eta^{\varepsilon'}). \quad (120)
\end{aligned}$$

This concludes the proof of the theorem. \square

9.2.2. *Proof of Theorem 6.1.1.* Recall that we want to prove the following result.

Theorem 9.2.2. *Let γ and η be two geodesics with at most one intersection point, then we have*

$$\lim_{n \rightarrow \infty} I_S(\{\hat{\ell}_{\gamma^n}(y), \hat{\ell}_{\eta^n}(y)\}) = \{\ell_\gamma, \ell_\eta\}_S$$

Proof. This will be a consequence of the generalized Wolpert Formula. By definition,

$$\hat{\ell}_\gamma(y) = \log(\mathbf{b}(\gamma^+, \gamma^-, \gamma^{-1}(y), \gamma(y))).$$

Thus

$$\{\hat{\ell}_\alpha(y), \hat{\ell}_\beta(y)\} = \sum_{\substack{u, u' \in \{-1, 1\} \\ v, v' \in \{-1, 1\}}} u \cdot u' \frac{\{\alpha^v \alpha^{-uv}(y), \beta^{v'} \beta^{-u'v'}(y)\}}{(\alpha^v \alpha^{-uv}(y)) \cdot (\beta^{v'} \beta^{-u'v'}(y))}. \quad (121)$$

But,

$$\begin{aligned}
& \{\alpha^v \alpha^{-uv}(y), \beta^{v'} \beta^{-u'v'}(y)\} \\
&= [(\alpha^v \alpha^{-uv}(y)), (\beta^{v'} \beta^{-u'v'}(y))] \alpha^v \beta^{-u'v'}(y) \cdot \beta^{v'} \alpha^{-uv}(y). \quad (122)
\end{aligned}$$

We remark that for n large enough, for all u, v, u', v' , we have

$$\begin{aligned} [(\gamma^v \gamma^{v.n}(y)), (\eta^{v'} \eta^{-u'.v'.n}(y))] &= 0, \\ [(\gamma^v \gamma^{-uv.n}(y)), (\eta^{v'} \eta^{v'.n}(y))] &= 0, \\ [(\gamma^v \gamma^{-v.n}(y)), (\beta^{v'} \eta^{-v.n}(y))] &= vv'[\gamma^+ \gamma^-, \eta^+ \eta^-]. \end{aligned} \quad (123)$$

Combining the remark in Equation (123), with (122) and (121), we have that for n large enough,

$$\{\hat{\ell}_{\gamma^n}(y), \hat{\ell}_\eta(y^n)\} = [\gamma^+ \gamma^-, \eta^+ \eta^-] \sum_{v, v' \in \{-1, 1\}} v.v' \frac{\gamma^v \eta^{-v'.n}(y) \cdot \eta^{v'} \gamma^{-v}(y)}{(\gamma^v \gamma^{-v.n}(y)) \cdot (\eta^{v'} \eta^{-v'.n}(y))} \quad (124)$$

Thus, taking the limit when n goes to ∞ yields

$$\begin{aligned} \lim_{n \rightarrow \infty} \left(\mathbf{I}_S \{ \hat{\ell}_{\gamma^n}(y), \hat{\ell}_\eta(y^n) \} \right) &= [\gamma^+ \gamma^-, \eta^+ \eta^-] \sum_{v, v' \in \{-1, 1\}} v.v' \frac{\gamma^v \eta^{-v'} \cdot \eta^{v'} \gamma^{-v}}{\gamma^v \gamma^{-v} \cdot \eta^{v'} \eta^{-v'}} \\ &= [\gamma^+ \gamma^-, \eta^+ \eta^-] \sum_{v, v' \in \{-1, 1\}} v.v' \cdot \mathbf{T}(\gamma^v, \eta^{v'}). \end{aligned} \quad (125)$$

The result now follows from this last equation and the generalised Wolpert formula (116). \square

10. DRINFEL'D-SOKOLOV REDUCTION

The purpose of this section is to prove Theorem 10.7.2 which explain the relation of the multi fraction algebra with the Poisson structure on $\mathrm{PSL}_n(\mathbb{R})$ -opers.

We spend the first three paragraphs explaining the Poisson structure on $\mathrm{PSL}_n(\mathbb{R})$ -opers using the Drinfel'd-Sokolov reduction of the Poisson structure on connections on the circle. Although this is a classical construction (see [3], [23] and [10] and the original reference [4]), we take some time explaining the main steps in differential geometric terms, expanding the sketch of the construction given by Graeme Segal in [22].

Finally, we relate the swapping algebra and this Poisson structure in Theorem 10.7.2.

10.1. Opers and non slipping connections. In this paragraph, we recall the definition $\mathrm{PSL}_n(\mathbb{R})$ -opers and show that they can be interpreted as class of equivalence of “non slipping” connections on a bundle with a flag structure.

10.1.1. *Opers.*

Definition 10.1.1. [OPERS] A $\mathrm{PSL}_n(\mathbb{R})$ -oper is a n^{th} -order linear differential operator on the circle $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ of the form

$$D : \psi \mapsto \frac{d^n \psi}{dt^n} + q_2 \frac{d^{n-2} \psi}{dt^{n-2}} + \dots + q_n \psi, \quad (126)$$

where q_i are functions.

Observe that this definition of an oper requires the choice of a parametrisation of the circle. Otherwise the q_i would rather be i -order differentials.

We denote by $X_n(\mathbb{T})$ the space of $\mathrm{PSL}_n(\mathbb{R})$ -opers on \mathbb{T} . Every oper has a natural holonomy which reflect the fact that the solutions may not be periodic. We consider the space $X_n(\mathbb{T})^0$ of opers with trivial holonomy, that is those opers D for which all solutions of $D\psi = 0$ are periodic. A Poisson structure on $X_n(\mathbb{T})$, whose symplectic leaves are opers with the same holonomy, was discovered in the context of integrable systems and Korteweg-de Vries equations – for a precise account of the history, see Dickey [3]. Later, Drinfeld-Solokolov interpreted that structure in a more differential geometric way in [4] and we shall now retrace the steps of that construction.

10.1.2. *Non slipping connections.* Let K be the line bundle of $(-1/2)$ -densities over \mathbb{T} and $P := J^{n-1}(K^{n-1})$ be the rank n vector bundle of $(n-1)$ jets of sections of the bundles of $(-(n-1)/2)$ -densities.

Let F_p be the vector subbundle of P defined by

$$F_p := \{j^{n-1}\sigma \mid j^{n-p-1}\sigma = 0\}.$$

The family $\{F_p\}_{1 \leq p \leq n}$ is a filtration of P : we have $F_n = P$, $F_{p-1} \subset F_p$ and $\dim(F_p) = p$.

The vector bundle also carries a natural volume form P , since $\det(P)$ is isomorphic to \mathbb{R} . Finally, $F_p/F_{p-1} = K^{2p}$.

We say a family of sections $\{e_1, \dots, e_n\}$ of P is a *basis for the filtration* if for every integer p no greater than n , for every $x \in S^1$, $\{e_1(x), \dots, e_p(x)\}$ is a basis of the fiber of F_p at x .

Definition 10.1.2. [NON-SLIPPING CONNECTIONS] A connection ∇ on P is non slipping if it satisfies the following conditions

- $\nabla F_p \subset F_{p+1}$ for all p ,
- If α_p is the projection from F_{p+1} to F_{p+1}/F_p , then the map

$$(X, u) \rightarrow \alpha_p(\nabla_X(u)),$$

considered as a linear map from $K^2 \otimes F_p/F_{p-1} = K^{2(p+1)}$ to $F_{p+1}/F_{p-1} = K^{2(p+1)}$ is the identity.

We denote by D_0 the space of non-slipping connections on P . The first classical proposition is that

Proposition 10.1.3. *Let ∇ be a non slipping connection, then there exists a unique basis $\{e_1, \dots, e_n\}$ of determinant 1 for the filtration so that*

$$\begin{aligned} \nabla_{\partial_t} e_i &= -e_{i+1}, \text{ for } i \leq n-1, \\ \nabla_{\partial_t} e_n &\in F_{n-1}, \end{aligned} \quad (127)$$

where ∂_t is the canonical vector field on \mathbb{T} .

Observe here that the basis depends on the choice of a parametrisation of the circle. From this proposition, it follows that we can associate to a non slipping connection ∇ the differential operator $D = D^\nabla$ so that

$$\nabla_{\partial_t}^* \left(\sum_{i=1}^{i=n} \frac{d^{i-1}\psi}{dt^{i-1}} \omega_i \right) = (D\psi)\omega_n,$$

where ∇^* is the dual connection and $\{\omega_i\}_{1 \leq i \leq n}$ is the dual basis to the basis $\{e_i\}_{1 \leq i \leq n}$ associated to ∇ in the previous proposition. One easily check that

$$D\psi = \frac{d^n \psi}{dt^n} + q_2 \frac{d^{n-2} \psi}{dt^{n-2}} + \dots + q_n \psi$$

where the functions q_i are given by $q_i = \omega_{n-j+1}(\nabla_{\partial_t} e_n)$.

We define

- (1) the *flag gauge group* as the group \mathbf{N} of linear automorphism of the bundle P defined by

$$\mathbf{N} := \{A \in \Omega^0(\mathbb{T}, \text{End}(P)) \mid A(F_p) = F_p, A|_{F_p/F_{p-1}} = \text{Id}\},$$

- (2) the *Lie algebra* \mathfrak{n} of the flag gauge group as

$$\mathfrak{n} := \{A \in \Omega^0(\mathbb{T}, \text{End}(P)) \mid A(F_p) \subset F_{p-1}\}.$$

We now have

Proposition 10.1.4. *The map $\nabla \mapsto D^\nabla$ realise an identification between D_0/\mathbf{N} and $\mathbf{X}_n(\mathbb{T})$ and this identification preserves the holonomy.*

It is interesting now to observe that the definition of an oper as an element of D_0/\mathbf{N} does not depend of a parametrisation.

Proof. Let ∇ be a non slipping connection $\{e_i\}$ the basis obtained by the previous proposition and $\nabla' = n.\nabla$ be a connection in the \mathbf{N} -orbit of ∇ . By definition of \mathbf{N} , $\nabla' e_i = \nabla e_i + u_i$, with $u_i \in F_{i-1}$. The result follows \square

10.2. The Poisson structure on the space of connections. The purpose of Drinfel'd-Sokolov reduction is to identify the space of opers $\mathbf{X}_n(\mathbb{T}) = \mathbf{D}_0/\mathbf{N}$ as a symplectic quotient of the space of all connexions on \mathbb{T} by the group \mathbf{N} .

Again, we shall paraphrase Segal, and define in this paragraph, as a first step of the construction of Drinfel'd-Sokolov reduction, the classical construction of the Poisson structure on the space of connections.

In general, when we deal with a Fréchet space of sections of a bundle, we have to specify functionals that we deemed observables and for which we can compute a Poisson bracket. This is done by specifying a subspace of cotangent vectors and describing the Poisson tensor on that subspace. Observables are then functionals whose differentials belong to that specific subset. However, the Poisson bracket can be extended to more general pair of observables. Rather than describing a general formalism – for which we could refer to [3] – we explain the construction in the case of connections.

10.2.1. Connexions and central extensions. Let \mathbf{G} be the Gauge group of the vector bundle P . The choice of a trivialisation of P give rise an isomorphism of \mathbf{G} with the loop group of $\mathrm{PSL}_n(\mathbb{R})$. Then, we introduce the following definitions

- (1) the *Lie algebra* \mathfrak{g} of \mathbf{G} is $\Omega^0(\mathbb{T}, \mathrm{End}_0(P))$ where $\mathrm{End}_0(P)$ stands for the vector space of trace free endomorphisms of P , this Lie algebra is equipped naturally with a coadjoint action of \mathbf{G} ,
- (2) the *dual Lie algebra* \mathfrak{g}° of \mathbf{G} is $\Omega^1(\mathbb{T}, \mathrm{End}_0(P))$,
- (3) the *duality* is given by the non degenerate bilinear mapping from $\mathfrak{g} \times \mathfrak{g}^\circ$ defined by

$$\langle \alpha, \beta \rangle = \int_{\mathbb{T}} \mathrm{tr}(\alpha \cdot \beta). \quad (128)$$

Let us choose a connection ∇ on P . Let Ω_∇ be the 2-cocycle on \mathfrak{g} given by the following formula

$$\Omega_\nabla(\xi, \eta) = \int_{\mathbb{T}} \mathrm{tr}(\xi \nabla \eta).$$

If ∇ and ∇' are two connections on P , then

$$\Omega_\nabla(\xi, \eta) - \Omega_{\nabla'}(\xi, \eta) = \alpha([\xi, \eta]),$$

where

$$\alpha(\chi) = \int_{\mathbb{T}} \mathrm{tr}((\nabla - \nabla') \cdot \chi).$$

In particular the cohomology class of the cocycle Ω_∇ does not depend on the choice of ∇ . Let $\widehat{\mathbf{G}}$ – whose Lie algebra is $\widehat{\mathfrak{g}}$ – be the central extension of \mathbf{G} corresponding of to this cocycle, so that

$$0 \rightarrow \mathbb{R} \rightarrow \widehat{\mathfrak{g}} \xrightarrow{\pi} \mathfrak{g}.$$

As we noticed, every connection defines a splitting of this sequence, that is a way to write $\widehat{\mathfrak{g}}$ as $\mathbb{R} \oplus \mathfrak{g}$.

Dually, we consider the vector space $\widehat{\mathfrak{g}}^\circ$ defined by the exact sequence

$$0 \xrightarrow{i} \mathfrak{g}^\circ \rightarrow \widehat{\mathfrak{g}}^\circ \rightarrow \mathbb{R},$$

with the duality with $\widehat{\mathfrak{g}}$ so that $\langle \gamma, i(\beta) \rangle = \langle \pi(\gamma), \beta \rangle$

It follows that the space of \mathcal{D} all $\mathrm{PSL}_n(\mathbb{R})$ connexions on P can be embedded in the space of these splittings which is identified with the affine hyperplane \mathbf{D} in $\widehat{\mathfrak{g}}^\circ$ defined by

$$\mathbf{D} := \{\beta \in \widehat{\mathfrak{g}}^\circ \mid \langle Z, \beta \rangle = 1\},$$

where $Z \in \widehat{\mathfrak{g}}$ the generator of the center. The hyperplane \mathbf{D} has \mathfrak{g}° as a tangent space. Observe that the embedding $\mathcal{D} \rightarrow \mathbf{D}$ is equivariant under the affine action of $\Omega^1(\mathbb{T}, \mathrm{End}_0(P)) = \mathfrak{g}^\circ \subset \widehat{\mathfrak{g}}^\circ$ as well as the coadjoint action of \mathbf{G} itself. In particular, the above embedding is surjective and we now identify \mathbf{D} as the space of all $\mathrm{PSL}_n(\mathbb{R})$ -connections on P . The coadjoint orbits of \mathbf{G} on \mathbf{D} are those connexions with the same holonomy.

10.2.2. The Poisson structure. Since we are working in infinite dimension, we are only going to define the Poisson tensor on certain “cotangent vectors” to \mathbf{D} . In our context we consider the set $\mathbf{D}^\circ := \mathfrak{g} = \Omega^0(\mathbb{T}, \mathrm{End}_0(P))$ of cotangent vectors where the duality is given by Formula (128). Using these notation, the Poisson structure is described in the following way

Definition 10.2.1. [POISSON STRUCTURE FOR CONNECTIONS]

- The Hamiltonian mapping from \mathbf{D}° to \mathbf{D} at a connection ∇ is

$$H : \alpha \mapsto d^\nabla \alpha.$$

- The Poisson tensor on \mathbf{D}° at a connection ∇ is

$$\Pi_\nabla(\alpha, \beta) := \langle \alpha, H(\beta) \rangle = \int_{\mathbb{T}} \mathrm{tr}(\alpha \cdot d^\nabla \beta).$$

- We say a functional F is an observable if its differential $d_\nabla F$ belongs to \mathbf{D}° for all ∇ . The Poisson bracket of two observables is

$$\{f, g\} := \Pi(df, dg) = \langle df, H(dg) \rangle.$$

REMARKS:

- (1) The Poisson bracket can be defined for more general pair of functionals than observables. Observe first that the differential of functionals on a Frechet space of sections of bundles – for instance connections – are distributions. Thus we can define the Poisson bracket of a general differentiable functional with an observable. For the purpose of this paper, we shall say that two functionals f and g form an *acceptable pair of observables* if their derivatives df and dg are distributions with disjoint singular support, or equivalently if they can be written as

$$\begin{aligned} df &= F + f_0 \\ dg &= G + g_0, \end{aligned}$$

where F and G have disjoint support and f_0, g_0 are observables in the previous meaning. In this case, their Poisson bracket is defined as

$$\{f, g\}(\nabla) = \Pi(f_0, g_0) + \langle F, H(g_0) \rangle - \langle G, H(f_0) \rangle.$$

This Poisson bracket agrees with regularising procedures.

- (2) We further observe that if D_∇ is the space of connections with the same holonomy as ∇ – that is the coadjoint orbit of ∇ – then the tangent space of D_∇ at ∇ is the vector space of exact 1-forms $d^\nabla(\Omega^0(\mathbb{T}, \text{End}_0(P)))$, and moreover the Poisson tensor on D_∇ is dual to the symplectic form ω defined by

$$\omega(d^\nabla\alpha, d^\nabla\beta) := \int_{\mathbb{T}} \text{tr}(\alpha \cdot d^\nabla\beta).$$

Thus the symplectic leaves of this Poisson structure are connections with the same holonomy. One can furthermore check that this formalism agrees with what we expect from coadjoint orbits.

10.3. Drinfel'd-Sokolov reduction. We now describe the Drinfel'd-Sokolov reduction. In the first paragraph we describe more precisely the group that we are going to work with in order to perform the reduction.

10.3.1. Dual Lie algebras. Let \mathfrak{n} be the Lie algebra of \mathbf{N} as defined above. Let \mathfrak{u} be the subspace of \mathfrak{g}° given by

$$\mathfrak{u} := \{A \in \Omega^1(\mathbb{T}, \text{End}_0(P)) \mid A(F_p) \subset F_p\}.$$

We observe

Proposition 10.3.1. *We have*

$$\mathfrak{u} = \{A \in \widehat{\mathfrak{g}}^\circ \mid \langle \alpha, A \rangle = 0, \quad \forall \alpha \in \mathfrak{n}\}$$

Thus if

$$\mathfrak{n}^\circ := \Omega^1(\mathbb{T}, \text{End}_0(P))/\mathfrak{u},$$

we have a duality $\mathfrak{n}^\circ \times \mathfrak{n} \rightarrow \mathbb{R}$ given by the map

$$\langle \alpha, \beta \rangle := \int_{\mathbb{T}} \text{tr}(\alpha\beta).$$

We now give another description of \mathfrak{n}° more suitable for our purpose. Let us first consider the natural projections

$$\begin{aligned} \pi_p^+ : \text{Hom}(F_p, E/F_p) &\rightarrow \text{Hom}(F_p, E/F_{p+1}), \\ \pi_p^- : \text{Hom}(F_p, E/F_p) &\rightarrow \text{Hom}(F_{p-1}, E/F_p). \end{aligned} \quad (129)$$

Let

$$M := \left\{ (u_1, \dots, u_{n-1}) \in \bigoplus_{p=1}^{p=n-1} \text{Hom}(F_p, E/F_p) \mid \pi_p^+(u_p) = \pi_{p+1}^-(u_{p+1}) \right\}.$$

We leave the reader check the following

Proposition 10.3.2. *The map defined from \mathfrak{n}° to $\Omega^1(\mathbb{T}, M)$ by*

$$A \rightarrow (A|_{F_1}, \dots, A|_{F_{n-1}})$$

is an isomorphism.

10.3.2. Drinfel'd-Sokolov reduction. If ∇ is a connection, we define the *slippage* – denoted $\sigma(\nabla)$ – of ∇ as the element of $\Omega^1(\mathbb{T}, M) = \mathfrak{n}^\circ$ given by

$$(u_1, \dots, u_p),$$

where $u_p(X, v) = \alpha_p(\nabla_X v)$ and α_p is the projection from E to E/F_p .

We are now going to define a canonical section of $\Omega^1(\mathbb{T}, M)$. We have a natural embedding

$$i_p : \text{Hom}(F_p/F_{p-1}, F_{p+1}/F_p) \rightarrow \text{Hom}(F_p, F/F_p).$$

Now observe that

$$\Omega^1(\mathbb{T}, \text{Hom}(F_p/F_{p-1}, F_{p+1}/F_p)) = (K^2)^* \otimes (K^{2p})^* \otimes K^{2(p+1)}.$$

Thus, let

$$\mathbf{I}_p := i_p(\text{Id}) \in (K^2)^* \otimes (K^{2p})^* \otimes K^{2(p+1)}.$$

Finally, we set

$$\mathbf{I} := (\mathbf{I}_1, \dots, \mathbf{I}_{n-1}),$$

and we observe that \mathbf{I} is invariant under the coadjoint action of \mathbf{N} .

Theorem 10.3.3. [DRINFEL'D-SOKOLOV REDUCTION] *The map σ is a moment map for the action of \mathbf{N} . Moreover $\mathbf{D}_0 = \mu^{-1}(\mathbf{I})$ and we thus obtain a Poisson structure on $\mathbf{X}_n(\mathbb{T})$.*

As a particular case of symplectic reduction, we briefly explain the construction of the Poisson bracket in our context of opers and non slipping connections. If f and g are two functionals on the space of opers, they are observables if their pull back F and G on the space of non slipping connections are observables and then their Poisson bracket is $\{f, g\}(D) := \{F, G\}(\nabla)$ where D is the oper associated with ∇ .

10.4. Opers and Frenet curves.

10.4.1. *Curves associated to $\mathrm{PSL}_n(\mathbb{R})$ -opers.* We recall that every oper D give rise to a curve from \mathbb{R} to $\mathbb{P}(\mathbb{R}^n)$ *equivariant* under the holonomy that is a curve

$$\xi : \mathbb{R} \rightarrow \mathbb{P}(\mathbb{R}^n),$$

such that $\xi(t+1) = H(\xi(t))$ where H is the holonomy. The construction runs as follows: the curve ξ is given in projective coordinates by

$$\xi := [v_1, \dots, v_n],$$

where $\{v_1, \dots, v_n\}$ are n -independent solutions of the equation $D\psi = 0$. The curve ξ is well defined up to the action of $\mathrm{PSL}_n(\mathbb{R})$. We call ξ the curve *associated* to the oper.

10.4.2. *Hitchin opers.* Let us say an oper is *Hitchin* if it has trivial holonomy and can be deformed through opers with trivial holonomy to the trivial oper $\psi \mapsto \frac{d^n \psi}{dt^n}$. Let us denote by $\mathbf{X}_n^0(\mathbb{T})$ the space of Hitchin opers which by the previous section inherits a Poisson structure.

10.4.3. *Frenet curves.* We say a curve ξ from \mathbb{T} to $\mathbb{P}(\mathbb{R}^n)$ is *Frenet* if there exists a curve $(\xi^1, \xi^2, \dots, \xi^{n-1})$ defined on \mathbb{T} , called the *osculating flag curve*, with values in the flag variety such that for every x in \mathbb{T} , $\xi(x) = \xi^1(x)$, and moreover

- For every pairwise distinct points (x_1, \dots, x_l) in \mathbb{T} and positive integers (n_1, \dots, n_l) such that

$$\sum_{i=1}^{i=l} n_i \leq n,$$

then the sum

$$\xi^{n_i}(x_i) + \dots + \xi^{n_l}(x_l)$$

is direct.

- For every x in \mathbb{T} and positive integers (n_1, \dots, n_l) such that

$$p = \sum_{i=1}^{i=l} n_i \leq n,$$

then

$$\lim_{\substack{(y_1, \dots, y_l) \rightarrow x, \\ y_i \text{ all distinct}}} \left(\bigoplus_{i=1}^{i=l} \xi^{n_i}(y_i) \right) = \xi^p(x).$$

We call $\xi^* := \xi^{n-1}$ the *osculating hyperplane*.

Since the trivial connection is non slipping with respect to the filtration given by osculating flags, we have the following obvious remark – see also [5] Section 9.12.

Proposition 10.4.1. *Every smooth Frenet curve comes from a $\mathrm{PSL}_n(\mathbb{R})$ -oper with trivial holonomy.*

Conversely, we now prove

Proposition 10.4.2. *The curve associated to a Hitchin oper is Frenet.*

Proof. Let us first introduce some notation and definitions. A *weighted* p -tuple X is a pair consisting of a p -tuple of pairwise distinct points (x_1, \dots, x_p) in \mathbb{T} – called the *support* – and a p -tuple of positive integers (j_1, \dots, j_p) so that $\sum_{1 \leq k \leq p} j_k = n$. If η is a curve defined on a subinterval I of \mathbb{T} with values in $\mathbb{R}^n \setminus \{0\}$, let

$$\hat{\eta}^{(p)}(x) := \eta(x) \wedge \dot{\eta}(x) \wedge \dots \wedge \eta^{(p-1)}(x) \in \Lambda^p(\mathbb{R}^n).$$

Moreover, if X is a weighted p -tuple as above with support in I , let

$$\hat{\eta}(X) := \bigwedge_{1 \leq k \leq p} \hat{\eta}^{(j_k)}(x^k) \in \Lambda^n(\mathbb{R}^n) = \mathbb{R}.$$

Say finally that a weighted p -tuple is *degenerate* with respect to η if $\hat{\eta}(X) = 0$. Observe finally that being degenerate only depends on the projection of η as a curve with values in $\mathbb{P}(\mathbb{R}^n)$ and thus makes sense for curves with values in $\mathbb{P}(\mathbb{R}^n)$. By definition, a curve ξ with values in $\mathbb{P}(\mathbb{R}^n)$ is Frenet if it admits no degenerate weighted p -tuple.

Let us work by contradiction and assume that there exists a Hitchin oper so that the associated curve is not Frenet. Let m be the smallest integer so that there exists a curve ξ associated to an Hitchin oper which admits a degenerate m -tuple.

Let \mathcal{O}_m be the set of Hitchin operators whose associated curve admits a degenerate m -tuple. By our standing assumption, \mathcal{O}_m is non empty, and moreover the trivial oper – which corresponds to the Veronese

embedding – does not belong to \mathcal{O}_m . Thus the contradiction will follow from the following discussion by the connexity of $X_n^0(\mathbb{T})$ where we prove that \mathcal{O}_m is both open and closed.

STEP 1: *The set \mathcal{O}_m is open in $X_n^0(\mathbb{T})$.*

Let

$$X = ((x^1, \dots, x^m), (i_1, \dots, i_m))$$

be a degenerate m -tuple for the curve ξ associated to the oper D . Without loss of generality, we can assume that i_1 is the greatest integer all integers j so that $((x^1, \dots, x^m), (j, \dots))$ is degenerate. Let now η be a lift (with values in $\mathbb{R}^n \setminus \{0\}$) of ξ on an interval containing the support of X . Let us consider the function f_D defined on a neighbourhood of x^1 by

$$f_D : y \mapsto \hat{\eta}(X(y)),$$

where $X(y) := ((y, x^2, \dots, x^m), (i_1, \dots, i_m))$. We first prove that $\dot{f}_D(x^1) \neq 0$. A computation yields

$$\dot{f}_D(x^1) = (\hat{\eta}^{(i_1-2)}(x_1) \wedge \eta^{(i_1)}(x_1)) \wedge \left(\bigwedge_{2 \leq j \leq m} \hat{\eta}^{i_j}(x_j) \right).$$

Let us recall the following elementary fact of linear algebra. let u, v and e_1, \dots, e_k are vectors in \mathbb{R}^n so that

$$u \wedge v \wedge e_1 \wedge \dots \wedge e_{k-1} \neq 0, \quad (130)$$

$$u \wedge e_1 \wedge \dots \wedge e_k = 0, \quad (131)$$

then

$$v \wedge e_1 \wedge \dots \wedge e_k \neq 0, \quad (132)$$

Indeed, by Assertion (131), u belongs to the hyperplane H generated by (e_1, \dots, e_k) . If Assertion (132) does not hold, then v also belongs to H . Thus the vector space plane generated by $(u, v, e_1, \dots, e_{k-1})$ also would lie in H , thus contradicting Assertion (130).

We know that by maximality of i_1 , $\hat{\eta}(Y) \neq 0$, where

$$Y = ((x^1, \dots, x^m), (i_1 + 1, i_2 - 1, \dots, i_m)) \neq 0.$$

Since $f_D(x^1) = 0$, the previous remark, with $u = \eta^{(i_1-1)}$, $v = \eta^{(i_1)}$ yields that $\dot{f}_D(x^1) \neq 0$.

By transversality, it then that for D' close to D there exist z close to x^1 so that $f_{D'}(z) = 0$ and thus $D' \in \mathcal{O}_m$.

STEP 2: *The set \mathcal{O}_m is closed in $X_n^0(\mathbb{T})$.*

Let $\{\xi_n^1\}_{n \in \mathbb{N}}$ be a sequence of curves associated to a converging sequence of opers in \mathcal{O}_m converging to an oper D associated to the curve ξ . Let

$$\{X_n = ((x_n^1, \dots, x_n^m), (j_n^1, \dots, j_n^m))\}_{n \in \mathbb{N}}.$$

be the corresponding sequence of degenerate m -tuples. We can extract a subsequence so that for every i , the sequence $\{j_n^i\}_{n \in \mathbb{N}}$ is constant and equal to j^i . After permutation of $\{1, \dots, p\}$ and extracting a further subsequence, we can assume that there exists a p -tuple, with $p \leq m$,

$$Y = ((y^1, \dots, y^p), (i^1, \dots, i^p)),$$

integers k_1, \dots, k_p so that

- (1) $1 = k_1 \leq \dots \leq k_p = m$
- (2) for all i , with $k_u \leq i < k_{u+1}$,

$$\lim_{n \rightarrow \infty} (x_n^i) = y^u,$$

- (3) for all v , with $1 \leq v \leq p$,

$$i^v = \sum_{k_v \leq u < k_{v+1}} j^u.$$

As an application of the Taylor formula, we have

$$\hat{\eta}^{(p)}(x) \wedge \hat{\eta}^{(k)}(y) = (x - y)^{p \cdot k} \hat{\eta}^{p+k}(x) + o((x - y)^k),$$

it follows that for all u

$$\lim_{n \rightarrow \infty} \left(\left(\prod_{v=k_u}^{k_{u+1}-1} \frac{1}{(x_n^v - y^u)^{N_v}} \right) \bigwedge_{v=k_u}^{k_{u+1}-1} \hat{\eta}^{(i^v)}(x_n^v) \right) = \hat{\eta}^{(i_u)}(y^u),$$

where $N_v = i^v (\sum_{w=k_u}^{v-1} i^w)$. In particular, Y is degenerate for ξ . Thus $p = m$ by minimality and $D \in \mathcal{O}_m$. □

Finally, let us say a Frenet curve is *Hitchin* if it can be deformed through Frenet curves to the Veronese embedding. Then we obtain a consequence of the two previous propositions the following statement which seems to belong to the folklore but for which we could not find a proper reference.

Theorem 10.4.3. *The map which associates to an $\mathrm{PSL}_n(\mathbb{R})$ -oper its associated curve is a homeomorphism from the space of Hitchin opers to the space of Hitchin Frenet curves.*

10.5. Cross ratios and opers. Let ξ be a Frenet curve and ξ^* be its associated osculating hyperplane curve. The *weak cross ratio* associated to this pair of curves is the function on

$$\mathbb{T}^{4*} := \{(x, y, z, t) \in \mathbb{T}^4 \mid z \neq y, x \neq t\},$$

defined by

$$\mathbf{b}_{\xi, \xi^*}(x, y, z, t) = \frac{\langle \widehat{\xi}(x) | \widehat{\xi}^*(y) \rangle \langle \widehat{\xi}(z) | \widehat{\xi}^*(t) \rangle}{\langle \widehat{\xi}(z) | \widehat{\xi}^*(y) \rangle \langle \widehat{\xi}(x) | \widehat{\xi}^*(t) \rangle}, \quad (133)$$

where for every u , we choose an arbitrary nonzero vector $\widehat{\xi}(u)$ and $\widehat{\xi}^*(u)$ respectively in $\xi(u)$ and $\xi^*(u)$. This weak cross ratio only depends on the oper D and we shall denote it by \mathbf{b}_D .

10.5.1. Coordinate functions. Let as in Section 10.1.2, K be the line bundle of $(-1/2)$ -densities over \mathbb{T} and $P := J^{n-1}(K^{n-1})$ be the rank n vector bundle of $(n-1)$ jets of sections of the bundle of $(-(n-1)/2)$ -densities. We choose once and for all a trivialisation of P given by n fibrewise independent sections $\sigma_1, \dots, \sigma_n$ of P so that F_p is generated by $\sigma_1, \dots, \sigma_p$.

Let ∇ be a connection on P . Let I be an interval in \mathbb{R} with extremities Y and y . We pull back ∇ , P and σ_i on \mathbb{R} using the projection

$$\pi : \mathbb{R} \rightarrow \mathbb{R}/\mathbb{Z} = \mathbb{T}.$$

We denote the pulled back objects by the same symbol. For any $y \in \mathbb{R}$, let σ_y the ∇ -parallel section of P on I characterised by $\sigma_y(y) = \sigma_1(y)$. Similarly let σ_Y^* be the ∇^* parallel section on I of P^* characterised by $\sigma_Y^*(Y) = \sigma_n^*(Y)$ where $(\sigma_1^*, \dots, \sigma_n^*)$ is the dual basis to $(\sigma_1, \dots, \sigma_n)$.

Then the function $t \mapsto \langle \sigma_Y^*(t), \sigma_y(t) \rangle$ is constant on I .

Definition 10.5.1. [COORDINATE FUNCTION] *The coordinate function associated to the points Y and y and the trivialization of P , is the function*

$$F_{Y,y} : \nabla \mapsto F_{Y,y}(\nabla) = \langle \sigma_Y^*(t), \sigma_y(t) \rangle,$$

defined on the space of connections on P .

We shall write $\sigma_Y^* \otimes \sigma_y =: \mathbf{p}^{Y,y} = \mathbf{p}^{Y,y}(\nabla) \in \Omega^0(\mathbb{R}, \text{End}_0(P))$ so that

$$F_{Y,y}(\nabla) = \text{tr}(\mathbf{p}^{Y,y}) \quad (134)$$

We then have

Proposition 10.5.2. *Assume that ∇ has trivial holonomy. Then the coordinate function $F_{Y,y}$ only depends on the projections of Y and y in \mathbb{T} . Moreover there exists a section $\mathbf{p}^{Y_0, y_0} \in \Omega^0(\mathbb{T}, \text{End}_0(P))$, so that $\mathbf{p}^{Y,y}$ is the pull back of \mathbf{p}^{Y_0, y_0} .*

Proof. Let Y_0 and y_0 the projections of Y and y , since ∇ has trivial holonomy, we may find parallel sections η_{y_0} and $\eta_{Y_0}^*$, so that $\eta_{y_0}(y_0) = \sigma_1(y_0)$ and $\eta_{Y_0}^*(Y_0) = \sigma_1(Y_0)$. Then, $\sigma_y = \pi^*(\eta_{y_0})$ and $\sigma_Y^* = \pi^*(\eta_{Y_0})$. Thus $F_{Y,y}(\nabla) = \langle \eta_{Y_0}^*(t), \eta_{y_0}(t) \rangle$. The first part of result follows. For the second part, we take $\mathbf{p}^{Y_0, y_0} = \eta_{Y_0}^* \otimes \eta_{y_0}$. \square

10.5.2. *Differential of coordinate functions.* Our aim in that paragraph is to compute the differential of $F_{Y,y}$, where Y, y belong to an interval I .

Proposition 10.5.3. *Let ∇ be a connection. Let y_0 be a point in $\mathbb{R} \setminus I$. Let α be an element of $\Omega^1(\mathbb{T}, \text{End}_0(P))$. Then*

$$\langle d_\nabla F_{Y,y}, \alpha \rangle = \int_{\mathbb{R}} \psi^{Y,y,y_0} \text{tr}(\mathbf{p}^{Y,y} \pi^*(\alpha)), \quad (135)$$

where $\psi^{Y,y,y_0}(s) := [y_0 s, Y y]$.

We can observe that the right hand side of Equation (135) does not depend on the choice of $y_0 \in \mathbb{R} \setminus I$. Indeed, by the cocycle identity, $\psi^{Y,y,x} - \psi^{Y,y,z}$ is constant and equal to $[xz, Y y] = 0$, if $x, z \notin I$.

Proof. Let β be a primitive of $\pi^* \alpha$ on I such that $\beta(y) = 0$. Let $t \mapsto \nabla^t$ be a one parameter smooth family of connections with $\nabla^0 = \nabla$ so that

$$\left. \frac{d}{dt} \right|_{t=0} \nabla^t = \alpha.$$

Let G^t be the family of sections of $\text{End}(P)$ so that $G^t(z) = \text{Id}$ and $(G^t)^* \nabla = \nabla^t$. Then by construction

$$\left. \frac{d}{dt} \right|_{t=0} G^t = \beta.$$

Moreover

$$F_{Y,y}(\nabla^t) = \langle \sigma_Y^*(Y), G^t(\sigma_y(Y)) \rangle.$$

Thus

$$\langle d_\nabla F_{Y,y}, \alpha \rangle = \langle \sigma_Y^*(Y), \beta(Y) \sigma_y(Y) \rangle. \quad (136)$$

Let $c(t)$ is a curve with value in I so that $c(0) = y$ and $c(1) = Y$. Let

$$f(t) = \langle \sigma_Y^*(c(t)), \beta(c(t)) \sigma_y^*(c(t)) \rangle.$$

Then

$$\begin{aligned} \langle d_\nabla F_{Y,y}, \alpha \rangle &= f(1) - f(0) \\ &= \int_0^1 \dot{f}(s) ds \end{aligned} \quad (137)$$

Since σ_Y^* and σ_y are parallel,

$$\dot{f}(s) = \sigma_Y^*(c(s)), \pi^*(\alpha(\dot{c}(s)).\sigma_y(c(s))),$$

and we have, letting J be the nerval whose endpoints are Y and y

$$\begin{aligned} \langle d_\nabla F_{Y,y}, \alpha \rangle &= \text{Sign}(Y - y) \int_J \langle \sigma_Y^*, \pi^*(\alpha).\sigma_y \rangle \\ &= \text{Sign}(Y - y) \int_J \text{tr}(\mathbf{p}^{Y,y}.\pi^*(\alpha)). \end{aligned} \quad (138)$$

We deduce the result from Equation (138) and the fact that if for any $y_0 \notin I$, we have

$$\text{Sign}(Y - y) \int_J \gamma = \int_{\mathbb{R}} \psi^{Y,y,y_0} \gamma.$$

□

10.6. Poisson brackets on the space of connections. Since $F_{X,x}$ is not an observable in the sense of Paragraph 10.2.2, we first need to regularise these functions

10.6.1. Regularisation. Let μ and ν two C^∞ measures compactly supported in a bounded interval $]a, b[$ of \mathbb{R} . Let us consider the function

$$F_{\mu,\nu} := \int_{\mathbb{R}^2} F_{X,x} d\mu.d\nu(X, x).$$

We consider this function defined on the space of connections over the bundle $P \rightarrow \mathbb{T}$. We obviously have

Proposition 10.6.1. *Let $\{(\mu_n, \nu_n)\}_{n \in \mathbb{N}}$ be two sequence of measures weakly converging to (μ, ν) , then $\{F_{\mu_n, \nu_n}\}_{n \in \mathbb{N}}$ converges uniformly on every compact to $F_{\mu, \nu}$.*

We say the sequence $\{(\mu_n, \nu_n)\}_{n \in \mathbb{N}}$ is *regularising* for the pair (X, x) if μ_n, ν_n are smooths measures weakly converging to the Dirac measures supported at X and x respectively.

10.6.2. Poisson brackets of regularisation. We now have

Proposition 10.6.2. *For any pair of smooth measures (μ, ν) with compact support, $F_{\mu,\nu}$ is an observable. Let (μ, ν) and $(\bar{\mu}, \bar{\nu})$ be two pairs of C^∞ measures on \mathbb{R} . Then the Poisson bracket $\{F_{\mu,\nu}, F_{\bar{\mu},\bar{\nu}}\}$ is equal to*

$$\sum_{m \in \mathbb{Z}} \int_{\mathbb{R}^4} [m(Y)m(y), Xx] \left(F_{X,y} F_{Y,x} - \frac{1}{n^2} F_{X,x} F_{Y,y} \right) d\Lambda(X, x, Y, y) \quad (139)$$

where $m(u) = u + m$ and $\Lambda = \mu \otimes \nu \otimes \bar{\mu} \otimes \bar{\nu}$. In particular if all measures are supported on $[0, 1]$, then the bracket $\{F_{\mu,\nu}, F_{\bar{\mu},\bar{\nu}}\}$ is equal to

$$\int_{[0,1]^4} [Yy, Xx] \left(F_{X,y} F_{Y,x} - \frac{1}{n^2} F_{X,x} F_{Y,y} \right) d\Lambda(X, x, Y, y). \quad (140)$$

Proof. By Proposition 10.5.3, we have that if a does not belong to the union K of the supports of μ and ν

$$\langle dF_{\mu,\nu}, \alpha \rangle = \int_{\mathbb{R}^2} \psi^{X,x,a} \operatorname{tr}(\mathfrak{p}^{X,x} \pi^* \alpha) d\mu.d\nu(X, x).$$

Let us denote by $C_0 := C - \frac{1}{n} \operatorname{tr}(C) \operatorname{Id}$ the trace free part of the endomorphism C . For any s in \mathbb{R} , let

$$\Lambda_{\mu,\nu}(s) := \int_{\mathbb{R}^2} \psi^{X,x,a}(s) \mathfrak{p}_0^{X,x}(s) d\mu.d\nu(X, x). \quad (141)$$

Observe that $\Lambda_{\mu,\nu} \in \Omega^0(\mathbb{R}, \operatorname{End}_0(P))$ and the support of $\Lambda_{\mu,\nu}$ is included in K . Let us trivialise P using the connection ∇ . Then let

$$G_{\mu,\nu}(s) := \sum_{m \in \mathbb{Z}} \Lambda_{\mu,\nu}(s + m).$$

Then $G_{\mu,\nu}(s)$ is periodic and thus of the form $\pi^* \beta$, with $\beta \in \Omega^0(\mathbb{T}, P)$. Moreover

$$\begin{aligned} \int_{\mathbb{T}} \operatorname{tr}(\beta.\alpha) &= \int_0^1 \operatorname{tr}(\pi^* \beta.\pi^* \alpha) \\ &= \int_{\mathbb{R}} \operatorname{tr}(\Lambda_{\mu,\nu}.\pi^* \alpha) \\ &= \langle dF_{\mu,\nu}, \alpha \rangle. \end{aligned} \quad (142)$$

It follows by Equation (142), that

$$dF_{\mu,\nu}(s) = \beta \in \mathfrak{g} = D^0. \quad (143)$$

In particular, according to Definition 10.2.1, $F_{\mu,\nu}$ is an observable. From Equation (141), we have

$$\begin{aligned} \Lambda_{\mu,\nu}(s) &= - \int_{-\infty}^s \int_s^\infty \mathfrak{p}_0^{X,x}(s) d\mu.d\nu(X, x) \\ &\quad + \int_s^\infty \int_{-\infty}^s \mathfrak{p}_0^{X,x}(s) d\mu.d\nu(X, x) \end{aligned}$$

For any smooth measure probability measure ξ let us write $d\xi = \dot{\xi} d\lambda$ where λ is the Lebesgue measure. Then, since $\mathfrak{p}^{X,x}$ is parallel, we

have

$$\begin{aligned}
\nabla_{\partial_t} \Lambda_{\mu,\nu}(s) &= -\dot{\mu}(s) \int_s^\infty \mathbf{p}_0^{s,x}(s) \, d\nu(x) + \dot{\nu}(s) \int_{-\infty}^s \mathbf{p}_0^{X,s}(s) \, d\mu(X) \\
&\quad -\dot{\mu}(s) \int_{-\infty}^s \mathbf{p}_0^{s,x}(s) \, d\nu(x) + \dot{\nu}(s) \int_s^\infty \mathbf{p}_0^{X,s}(s) \, d\mu(X) \\
&= \dot{\nu}(s) \int_{\mathbb{R}} \mathbf{p}_0^{X,s} d\mu(X) - \dot{\mu}(s) \int_{\mathbb{R}} \mathbf{p}_0^{s,x} d\nu(x).
\end{aligned}$$

It follows that

$$\begin{aligned}
&\text{tr}(\Lambda_{\mu,\nu}(s+m) \nabla_{\partial_t} \Lambda_{\bar{\mu},\bar{\nu}}(s)) \\
&= \dot{\nu}(s) \int_{\mathbb{R}^3} \psi^{X,x,a}(s+m) \text{tr}(\mathbf{p}_0^{Y,s} \mathbf{p}_0^{X,x}) \, d\mu.d\nu.d\bar{\mu}(X,x,Y) \\
&\quad - \dot{\mu}(s) \int_{\mathbb{R}^3} \psi^{X,x,a}(s+m) \text{tr}(\mathbf{p}_0^{s,y} \mathbf{p}_0^{X,x}) \, d\mu.d\nu.d\bar{\nu}(X,x,y) \quad (144)
\end{aligned}$$

We can now compute the Poisson bracket as defined in Definition 10.2.1:

$$\begin{aligned}
\{F_{\mu,\nu}, F_{\bar{\mu},\bar{\nu}}\} &= \int_{\mathbb{R}} \text{tr}(\Lambda_{\mu,\nu}(s) \pi^*(\nabla_{\partial_t} dF_{\bar{\mu},\bar{\nu}}(s))) \, d\lambda(s) \\
&= \sum_{m \in \mathbb{Z}} \int_{\mathbb{R}} \text{tr}(\Lambda_{\mu,\nu}(s) \nabla_{\partial_t} \Lambda_{\bar{\mu},\bar{\nu}}(s+m)) \, d\lambda(s) \\
&= \sum_{m \in \mathbb{Z}} \int_{\mathbb{R}} \text{tr}(\Lambda_{\mu,\nu}(s+m) \nabla_{\partial_t} \Lambda_{\bar{\mu},\bar{\nu}}(s)) \, d\lambda(s) \quad (145)
\end{aligned}$$

Using Equation (144), we get that

$$\begin{aligned}
&\{F_{\mu,\nu}, F_{\bar{\mu},\bar{\nu}}\} \\
&= \sum_{m \in \mathbb{Z}} \int_{\mathbb{R}^4} \psi^{X,x,a}(s+m) \text{tr}(\mathbf{p}_0^{Y,s} \mathbf{p}_0^{X,x}) \, d\Lambda(X,x,Y,s) \quad (146)
\end{aligned}$$

$$- \sum_{m \in \mathbb{Z}} \int_{\mathbb{R}^4} \psi^{X,x,a}(s+m) \text{tr}(\mathbf{p}_0^{s,y} \mathbf{p}_0^{X,x}) \, d\Lambda(X,x,s,y) \quad (147)$$

Using the dummy changes of variables $s = y$ on Line (146) and $s = Y$ on Line (147), we finally get

$$\begin{aligned}
&\{F_{\mu,\nu}, F_{\bar{\mu},\bar{\nu}}\} \\
&= \sum_{m \in \mathbb{Z}} \int_{\mathbb{R}^4} (\psi^{X,x,a}(y+m) - \psi^{X,x,a}(Y+m)) \text{tr}(\mathbf{p}_0^{Y,y} \mathbf{p}_0^{X,x}) \, d\Lambda(X,x,Y,y) \\
&= \sum_{m \in \mathbb{Z}} \int_{\mathbb{R}^4} [(Y+m)(y+m), Xx] \text{tr}(\mathbf{p}_0^{Y,y} \mathbf{p}_0^{X,x}) \, d\Lambda(X,x,Y,y) \quad (148)
\end{aligned}$$

We conclude the proof of the proposition by remarking that

$$\mathrm{tr}(\mathbf{p}^{X,x} \mathbf{p}^{Y,y}) = \mathrm{tr}(\mathbf{p}^{X,y}) \mathrm{tr}(\mathbf{p}^{Y,x}),$$

and thus

$$\mathrm{tr}(\mathbf{p}_0^{X,x} \mathbf{p}_0^{Y,y}) = \mathrm{tr}(\mathbf{p}^{X,y}) \mathrm{tr}(\mathbf{p}^{Y,x}) - \frac{1}{n^2} \mathrm{tr}(\mathbf{p}^{X,x}) \mathrm{tr}(\mathbf{p}^{Y,y}). \quad (149)$$

Combining Equations (148) and (149) yields the result. \square

As corollaries, we obtain

Corollary 10.6.3. *Let (μ_u, ν_n) and $(\bar{\mu}_n, \bar{\nu}_n)$ be regularising sequences for (X, x) and (Y, y) respectively. Assume that $\{X, x, Y, y\} \subset]0, 1[$, then*

$$\lim_{n \rightarrow \infty} (\{F_{\mu_n, \nu_n}, F_{\bar{\mu}_n, \bar{\nu}_n}\}) = [Yy, Xx] \left(F_{X,y} F_{Y,x} - \frac{1}{n^2} F_{X,x} F_{Y,y} \right).$$

Corollary 10.6.4. *Let (X, x, Y, y) be a quadruple of pairwise distinct points. Then $(F_{X,x}, F_{Y,y})$ is a pair of acceptable observables. Moreover,*

$$\{F_{X,x}, F_{Y,y}\} = [Yy, Xx] \left(F_{X,y} F_{Y,x} - \frac{1}{n^2} F_{X,x} F_{Y,y} \right).$$

This last corollary interprets the swapping algebra as an algebra of “observables” on the space of connections.

10.7. Drinfel’d-Sokolov reduction and the multi fraction algebra. We intruced in Paragraph 10.5.1 functions of connections depending on the choice of a trivialisation of P . We now introduce functions that only depend on the associated oper and do not rely on the choice of the trivialisation of P .

We first relate crossratios with our previously defined coordinate functions.

10.7.1. Cross ratios. The following proposition follows at once from the definitions.

Proposition 10.7.1. *Let D a Hitchin oper associated to connection ∇ with trivial holonomy. Let X, x, Y, y be a quadruple of pairwise distinct points of \mathbb{T} . Let $\tilde{X}, \tilde{x}, \tilde{Y}, \tilde{y}$ be lifts of X, x, Y, y in \mathbb{R} , then*

$$\mathbf{b}_D(X, x, Y, y) = \frac{F_{\tilde{X}, \tilde{y}}(\nabla) \cdot F_{\tilde{Y}, \tilde{x}}(\nabla)}{F_{\tilde{X}, \tilde{x}}(\nabla) \cdot F_{\tilde{Y}, \tilde{y}}(\nabla)}.$$

10.7.2. *The main Theorem.* We can now prove our main theorem which relates the Poisson structure on the space of opers and the multi fraction algebra.

Theorem 10.7.2. *Let $(X_0, x_0, Y_0, y_0, X_1, x_1, Y_1, y_1)$ be pairwise distinct points. Then the cross fractions $[X_0; x_0; Y_0; y_0]$ and $[X_1; x_1; Y_1; y_1]$ defines a pair of acceptable observables whose Poisson bracket with respect to the Drinfel'd-Sokolov reduction coincide with their Poisson bracket in the multi fraction algebra.*

Proof. This is an immediate consequence of Proposition 10.7.1 and Corollary 10.6.4, as well as the definition of the Poisson structure coming from the symplectic reduction in Theorem 10.3.3. \square

11. APPENDIX: EXISTENCE OF VANISHING SEQUENCES

We prove the existence of vanishing sequences.

Definition 11.0.3. [SEPARABILITY IN GROUPS] *Let G be a group. The group G is said to be subgroup separable, if given any finitely generated subgroup H in G , given any $g \in G$ and $h \notin Hg$, there exists a finite index subgroup G_0 in G such that if π is the projection of G onto G/G_0 , then*

$$\pi(h) \notin \pi(Hg).$$

Let G be a group. The group G is said to be double coset separable, if given any finitely generated subgroups H and K in G , given any $g \in G$ and $h \notin HgK$, there exists a finite index subgroup G_0 in G such that if π is the projection of G onto G/G_0 , then

$$\pi(h) \notin \pi(HgK).$$

Observe that a double coset separable group is then subgroup separable and residually finite. G. Niblo proved in [19]

Theorem 11.0.4. *A surface group is double coset separable.*

As an immediate consequence, since $\pi_1(S)$ is countable, we have

Corollary 11.0.5. *Vanishing sequences exist.*

REFERENCES

- [1] Michael F Atiyah and Raoul Bott, *The Yang-Mills equations over Riemann surfaces*, Philos. Trans. Roy. Soc. London Ser. A **308** (1983), no. 1505, 523–615.
- [2] Marc Bourdon, *Sur le birapport au bord des $\text{CAT}(-1)$ -espaces*, Publ. Math. IHES (1996), no. 83, 95–104.
- [3] L. A Dickey, *Lectures on classical W -algebras*, Acta Appl. Math. **47** (1997), no. 3, 243–321.
- [4] V. G. Drinfel'd and V. V. Sokolov, *Equations of Korteweg-de Vries type, and simple Lie algebras*, Dokl. Akad. Nauk SSSR **258** (1981), no. 1, 11–16. MR 615463 (83k:58040)
- [5] Vladimir V Fock and Alexander B Goncharov, *Moduli spaces of local systems and higher Teichmüller theory*, Publ. Math. IHES (2006), no. 103, 1–211.
- [6] William M Goldman, *The symplectic nature of fundamental groups of surfaces*, Adv. Math. **54** (1984), no. 2, 200–225.
- [7] ———, *Invariant functions on Lie groups and Hamiltonian flows of surface group representations*, Invent. Math. **85** (1986), no. 2, 263–302.
- [8] Suresh Govindarajan, *Higher-dimensional uniformisation and W -geometry*, Nuclear Phys. B **457** (1995), no. 1-2, 357–374.
- [9] Suresh Govindarajan and T. Jayaraman, *A proposal for the geometry of W_n -gravity*, Phys. Lett. B **345** (1995), no. 3, 211–219.
- [10] Partha Guha, *Euler–Poincaré flows on SL_n -opers and integrability*, Acta Applicandae Mathematicae: An International Survey ... (2007).
- [11] Olivier Guichard, *Composantes de Hitchin et représentations hyperconvexes de groupes de surface*, J. Differential Geom (2008).
- [12] Nigel J Hitchin, *Lie groups and Teichmüller space*, Topology **31** (1992), no. 3, 449–473.
- [13] François Labourie, *Anosov flows, surface groups and curves in projective space*, Invent. Math. **165** (2006), no. 1, 51–114.
- [14] ———, *Cross ratios, surface groups, $\text{PSL}(n, \mathbf{R})$ and diffeomorphisms of the circle*, Publ. Math. IHES (2007), no. 106, 139–213.
- [15] ———, *Flat projective structures on surfaces and cubic holomorphic differentials*, Pure Appl. Math. Q. **3** (2007), no. 4, 1057–1099.
- [16] François Labourie, *An algebra of observables for cross ratios*, C. R. Math. Acad. Sci. Paris **348** (2010), no. 9-10, 503–507. MR 2645161 (2011h:17032)
- [17] François Ledrappier, *Structure au bord des variétés à courbure négative*, vol. 13, 1995, pp. 97–122.
- [18] Franco Magri, *A simple model of the integrable Hamiltonian equation*, J. Math. Phys. **19** (1978), no. 5, 1156–1162. MR 488516 (80a:35112)
- [19] G Niblo, *Separability properties of free groups and surface groups*, J. Pure Appl. Algebra **78** (1992), no. 1, 77–84.
- [20] Jean-Pierre Otal, *Le spectre marqué des longueurs des surfaces à courbure négative*, Ann. of Math. (2) **131** (1990), no. 1, 151–162.
- [21] ———, *Sur la géométrie symplectique de l'espace des géodésiques d'une variété à courbure négative*, Rev. Mat. Iberoamericana **8** (1992), no. 3, 441–456.
- [22] Graeme Segal, *The geometry of the KdV-equation*, Internat. J. Modern Phys. A **6** (1991), no. 16, 2859–2869.

- [23] Pierre van Moerbeke, *Algèbres W et équations non-linéaires*, Astérisque (1998), no. 252, Exp. No. 839, 3, 105–129.
- [24] E Witten, *Surprises with topological field*, cdsweb.cern.ch.
- [25] Scott A. Wolpert, *The Fenchel-Nielsen deformation*, Ann. of Math. (2) **115** (1982), no. 3, 501–528. MR 657237 (83g:32024)
- [26] ———, *On the symplectic geometry of deformations of a hyperbolic surface*, Ann. of Math. (2) **117** (1983), no. 2, 207–234.